

ACTEMRA SQ

Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS
- TYENNE AUTOINJECTOR
- TYENNE SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	Interstitial lung disease-18 years and older (initial and continuation)
Prescriber Restrictions	RA/GCA/PJIA/SJIA - Prescribed by or in consultation with a rheumatologist (initial therapy only). Lung disease-presc/consult-pulmonologist or rheum (initial and cont)
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: RHEUMATOID ARTHRITIS (RA) [A or B]: A) tried two of the following: Enbrel, preferred adalimumab product (see Example 1), Orencia, Rinvoq or Xeljanz/XR (Note: trials with the following will also count towards meeting the try two requirement: Cimzia, infliximab, Kevzara, golimumab SC/IV, non-preferred adalimumab product), or B) heart failure or a previously treated lymphoproliferative disorder. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA) [A or B]: A) tried two of the following: Enbrel, Orencia, Rinvoq, Xeljanz, preferred adalimumab product. (Note: trials with Kevzara, infliximab or a non-preferred adalimumab product will also count towards meeting the try two requirement), or B) heart failure or a previously treated lymphoproliferative disorder. SYSTEMIC-ONSET JIA (SJIA): Approve. GIANT CELL ARTERITIS: tried or is currently taking a systemic CS or CS is contraindicated. INTERSTITIAL LUNG DISEASE ASSOCIATED WITH SYSTEMIC SCLEROSIS (A and B): A) elevated acute phase reactants and B) diagnosis confirmed by high-resolution computed tomography. CONTINUATION THERAPY: ALL INDICATIONS: patient had a response to therapy. Example 1: preferred adalimumab products include Humira (NDCs starting with -00074), Cyltezo, Yuflyma.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ACTIMMUNE

Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Chronic granulomatous disease - prescribed by or in consultation with an immunologist, hematologist or infectious disease specialist. Malignant osteopetrosis- prescribed by or in consultation with an endocrinologist or hematologist.
Coverage Duration	1 year
Other Criteria	Chronic granulomatous disease - approve if diagnosis has been established by a molecular genetic test identifying a gene-related mutation linked to chronic granulomatous disease. Malignant osteopetrosis, severe - approve if pt has had radiographic (X-ray) imaging demonstrating skeletal features related to osteopetrosis or pt had a molecular genetic test identifying a gene-related mutation linked to severe, malignant osteopetrosis.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ACYCLOVIR (TOPICAL)

Products Affected

- *acyclovir topical ointment*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ADALIMUMAB

Products Affected

- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML
- CYLTEZO(CF) PEN PSORIASIS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML
- CYLTEZO(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.2 ML, 20 MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8 ML
- HUMIRA (PREFERRED NDCS STARTING WITH 00074) SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA PEN (PREFERRED NDCS STARTING WITH 00074)
- HUMIRA(CF) (PREFERRED NDCS STARTING WITH 00074) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML
- HUMIRA(CF) PEN (PREFERRED NDCS NDCS STARTING WITH 00074) SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 80 MG/0.8 ML
- HUMIRA(CF) PEN CROHNS-UC-HS (PREFERRED NDCS NDCS STARTING WITH 00074)
- HUMIRA(CF) PEN PSOR-UV-ADOL HS (PREFERRED NDCS NDCS STARTING WITH 00074)
- YUFLYMA(CF) AI CROHN'S-UC-HS
- YUFLYMA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR, KIT 40 MG/0.4 ML, 80 MG/0.8 ML
- YUFLYMA(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another biologic DMARD or targeted synthetic DMARD.
Required Medical Information	Diagnosis
Age Restrictions	Pending CMS Review
Prescriber Restrictions	Pending CMS Review
Coverage Duration	Approve through end of plan year
Other Criteria	Pending CMS Review
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pending CMS Review

PA Criteria	Criteria Details
Part B Prerequisite	No

ADBRY

Products Affected

- ADBRY

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another monoclonal antibody therapy (i.e., Dupixent, Cinqair, Fasenra, Nucala, Tezspire, or Xolair). Concurrent use with Janus Kinase Inhibitors (JAKis) [oral or topical].
Required Medical Information	Diagnosis
Age Restrictions	AD-12 years of age and older (initial therapy)
Prescriber Restrictions	Atopic Dermatitis-prescribed by or in consultation with an allergist, immunologist or dermatologist (initial therapy)
Coverage Duration	Initial-Atopic Dermatitis-4 months, Continuation-1 year
Other Criteria	Atopic Dermatitis, initial-patient has atopic dermatitis involvement estimated to be greater than or equal to 10 percent of the body surface area and patient meets a and b: a. Patient has tried at least one medium-, medium-high, high-, and/or super-high-potency prescription topical corticosteroid AND b. Inadequate efficacy was demonstrated with the previously tried topical corticosteroid therapy. Continuation- Approve if the patient has been receiving Adbry for at least 4 months and patient has responded to therapy. Note: A patient who has received less than 4 months of therapy or who is restarting therapy with Adbry should be considered under initial therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with Phosphodiesterase Inhibitors Used for Pulmonary Hypertension or Other Soluble Guanylate Cyclase Stimulators.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	PAH and CTEPH- must be prescribed by or in consultation with a cardiologist or a pulmonologist.
Coverage Duration	1 year
Other Criteria	For PAH - must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AIMOVIG

Products Affected

- AIMOVIG AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Combination therapy with another cGRP inhibitor for migraine headache prevention
Required Medical Information	Diagnosis, number of migraine headaches per month
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Approve if the patient meets the following criteria (A and B): A) Patient has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication), AND B) If pt is currently taking Aimovig, the pt has had significant clinical benefit from the medication.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AKEEGA

Products Affected

- AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate cancer- Approve if the patient meets the following (A, B, C, and D): A)Patient has metastatic castration-resistant prostate cancer, AND B)Patient has a BReast CAncer (BRCA) mutation, AND C)The medication is used in combination with prednisone, AND D)Patient meets one of the following (i or ii): i. The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog, Note: Examples are leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix acetate subcutaneous injection), and Orgovyx (relugolix tablets).OR ii. Patient has had a bilateral orchiectomy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ALECENSA

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Pediatric diffuse high grade glioma- less than or equal to 21 years old, All others- 18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Non-small cell lung cancer-approve if the patient has both (A and B): A either (i or ii): i) medication is used as adjuvant treatment following tumor resection (note: for tumors greater than or equal to 4 cm or node positive) or ii) advanced or metastatic disease and B) anaplastic lymphoma kinase (ALK)-positive disease as detected by an approved test. Anaplastic large cell lymphoma-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease and (i or ii): (i) the medication is used for palliative-intent therapy, or (ii) pt has relapsed or refractory disease. Erdheim-Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory Myofibroblastic Tumor- pt has anaplastic lymphoma kinase (ALK)-positive disease AND (i or ii): (i) pt has advanced, recurrent or metastatic disease, or (ii) tumor is inoperable. Large B-Cell Lymphoma- pt has ALK-positive disease AND pt has relapsed or refractory disease. Pediatric diffuse high grade glioma- approve if (A and B): A) ALK-positive disease, and B) either (i or ii): i) medication is used as adjunctive treatment AND tumor is not diffuse midline glioma, H3 K27-altered or pontine location, or ii) medication is used for recurrent or progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Anaplastic large cell lymphoma, Erdheim Chester disease, Inflammatory Myofibroblastic Tumor, Large B-Cell Lymphoma, Pediatric Diffuse High Grade Glioma
Part B Prerequisite	No

ALOSETRON

Products Affected

- *alosetron*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ALPHA 1 PROTEINASE INHIBITORS

Products Affected

- PROLASTIN-C INTRAVENOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Alpha1-Antitrypsin Deficiency with Emphysema (or Chronic Obstructive Pulmonary Disease)-approve if the patient has a baseline (pretreatment) AAT serum concentration of less than 80 mg/dL or 11 micromol/L.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ALUNBRIG

Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	ALK status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Erdheim-Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory myofibroblastic tumor (IMT)-approve if the patient has ALK positive disease and has advanced, recurrent or metastatic disease or the tumor is inoperable. NSCLC, must be ALK-positive, as detected by an approved test, have advanced or metastatic disease and patients new to therapy must have a trial of Alecensa or Lorbrena prior to approval of Alunbrig. Peripheral T-Cell Lymphoma- approve if patient has ALK-positive anaplastic large cell lymphoma (ALCL).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Erdheim-Chester disease, Inflammatory myofibroblastic tumor (IMT), Peripheral T-Cell Lymphoma
Part B Prerequisite	No

ANTIBIOTICS (IV)

Products Affected

- *amikacin injection solution 500 mg/2 ml*
- *ampicillin sodium injection recon soln 1 gram, 10 gram*
- *ampicillin-sulbactam injection*
- *azithromycin intravenous*
- *aztreonam*
- *BICILLIN L-A*
- *cefoxitin*
- *ceftazidime*
- *cefuroxime sodium injection recon soln 750 mg*
- *cefuroxime sodium intravenous recon soln 1.5 gram*
- *ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml*
- *clindamycin in 5 % dextrose*
- *clindamycin phosphate injection*
- *colistin (colistimethate na)*
- *doxy-100*
- *doxycycline hyclate intravenous*
- *ertapenem*
- *gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml*
- *gentamicin injection*
- *imipenem-cilastatin*
- *levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml*
- *linezolid in dextrose 5%*
- *meropenem intravenous recon soln 1 gram, 500 mg*
- *metronidazole in nacl (iso-osm)*
- *moxifloxacin-sod.chloride(iso)*
- *nafcillin injection*
- *oxacillin*
- *oxacillin in dextrose(iso-osm) intravenous piggyback 2 gram/50 ml*
- **PENICILLIN G POT IN DEXTROSE INTRAVENOUS PIGGYBACK 2 MILLION UNIT/50 ML, 3 MILLION UNIT/50 ML**
- *penicillin g potassium injection recon soln 20 million unit*
- *penicillin g sodium*
- **STREPTOMYCIN**
- *tazicef injection*
- **TEFLARO**
- *tigecycline*
- *tobramycin sulfate injection solution*
- *vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A

PA Criteria	Criteria Details
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ANTIFUNGALS (IV)

Products Affected

- *fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml*
- *voriconazole*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ARCALYST

Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent biologic therapy
Required Medical Information	N/A
Age Restrictions	Initial tx CAPS/Pericarditis-Greater than or equal to 12 years of age.
Prescriber Restrictions	Initial tx CAPS-prescribed by, or in consultation with, a rheumatologist, geneticist, allergist/immunologist, or dermatologist. DIRA initial-rheum, geneticist, derm, or a physician specializing in the treatment of autoinflammatory disorders. Pericarditis-cardiologist or rheum
Coverage Duration	CAPS-3 mos initial, 1 yr cont. DIRA-6 mos initial, 1 yr cont. Pericard-3 mos initial, 1 yr cont
Other Criteria	INITIAL THERAPY: DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA) [all of A, B, and C]: A) weighs at least 10 kg, B) genetic test confirms a mutation in the IL1RN gene, and C) had clinical benefit with anakinra subcutaneous injection. PERICARDITIS: pericarditis is recurrent. CONTINUATION THERAPY: ALL INDICATIONS: patient had a positive response to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ARIKAYCE

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous medication history (as described in Other Criteria field)
Age Restrictions	MAC-18 years and older (initial therapy)
Prescriber Restrictions	MAC initial-Prescribed by a pulmonologist, infectious disease physician or a physician who specializes in the treatment of MAC lung infections.
Coverage Duration	1 year
Other Criteria	<p>INITIAL THERAPY: MYCOBACTERIUM AVIUM COMPLEX (MAC) LUNG DISEASE (all of A, B, and C): A) positive sputum culture for MAC [Note: any positive sputum culture taken after completion of a background multidrug regimen (throughout, see Example 1 below) fulfills this criterion], B) MAC isolate is susceptible to amikacin, and C) Arikayce will be used in combination with a background multidrug regimen.</p> <p>CONTINUATION THERAPY: MAC LUNG DISEASE (A and B): A) Arikayce prescribed in combination with a background multidrug regimen and B) patient meets one of the following (a or b): a) patient has not achieved negative sputum cultures for MAC or b) patient has achieved negative sputum cultures for MAC for less than 12 months. Example 1: background multidrug regimen example - a macrolide (azithromycin or clarithromycin), ethambutol, and a rifamycin (rifampin or rifabutin).</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AUBAGIO

Products Affected

- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of teriflunomide with other disease-modifying agents used for multiple sclerosis (MS)
Required Medical Information	Relapsing form of MS, to include, clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AUGTYRO

Products Affected

- AUGTYRO ORAL CAPSULE 160 MG,
40 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	NSCLC - 18 and older, Solid tumors - 12 and older, Pediatric Diffuse High-Grade Glioma-less than 18
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Non-Small Cell Lung Cancer-approve if the patient has locally advanced or metastatic disease, patient has ROS1-positive non-small cell lung cancer and the mutation was detected by an approved test. Note: If the patient has non-small cell lung cancer with neurotrophic receptor tyrosine kinase (NTRK) gene fusion, see Solid Tumors indication. Solid tumors - approve if tumor is positive for neurotrophic tyrosine receptor kinase (NTRK) gene fusion AND tumor is locally advanced or metastatic or surgical resection will likely result in severe morbidity. Pediatric Diffuse High-Grade Gliomas - approve if tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the medication is used as adjuvant therapy or for recurrent or progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pediatric Diffuse High-Grade Gliomas
Part B Prerequisite	No

AUSTEDO

Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 18 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	Chorea-prescribed by or in consult with a neuro. TD-Prescribed by or in consultation with a neurologist or a psychiatrist
Coverage Duration	1 year
Other Criteria	Chorea associated with Huntington's Disease-approve if the diagnosis of Huntington's Disease is confirmed by genetic testing. Tardive dyskinesia-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AVMAPKI-FAKZYNJA

Products Affected

- AVMAPKI-FAKZYNJA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER- ALL of the following (A, B and C): A) Patient has recurrent low-grade serous cancer, AND B) The cancer has a KRAS mutation, AND C) Patient has tried at least one systemic therapy. Note: Examples of systemic therapy include one or more of the following medications: paclitaxel, carboplatin, bevacizumab, letrozole, anastrozole, or exemestane.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Fallopian Tube or Primary Peritoneal Cancer
Part B Prerequisite	No

AVONEX

Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of other disease-modifying agent used for multiple sclerosis
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or after consultation with a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AYVAKIT

Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	GIST-approve if the tumor is positive for platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation or if the patient has tried two of the following: Gleevec (imatinib), Sutent (sunitinib), Sprycel (dasatinib), Stivarga (regorafenib) or Qinlock (ripretinib). Myeloid/Lymphoid Neoplasms with eosinophilia-approve if the tumor is positive for PDGFRA D842V mutation. Systemic mastocytosis-Approve if the patient has a platelet count greater than or equal to 50,000/mcL and patient has either indolent systemic mastocytosis or one of the following subtypes of advanced systemic mastocytosis-aggressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm or mast cell leukemia.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myeloid/Lymphoid neoplasms with Eosinophilia
Part B Prerequisite	No

BALVERSA

Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies, test results
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Urothelial Carcinoma, locally advanced or metastatic-approve if the patient has susceptible fibroblast growth factor receptor 3 genetic alterations AND the patient has progressed during or following prior platinum-containing chemotherapy, other chemotherapy or checkpoint inhibitor therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BENLYSTA

Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with Other Biologics or Lupkynis
Required Medical Information	Diagnosis
Age Restrictions	5 years and older (initial).
Prescriber Restrictions	SLE-Prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist or dermatologist (initial and continuation). Lupus Nephritis-nephrologist or rheum. (Initial/cont)
Coverage Duration	SLE-Initial-4 months, cont-1 year. Lupus Nephritis-6 mo initial, 1 year cont
Other Criteria	Lupus Nephritis Initial-approve if the patient has a diagnosis of lupus nephritis confirmed on biopsy (For example, World Health Organization class III, IV, or V lupus nephritis), AND the medication is being used concurrently with an immunosuppressive regimen (ex: azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate mofetil and/or a systemic corticosteroid). Cont-approve if the medication is being used concurrently with an immunosuppressive regimen (ex: azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate mofetil and/or a systemic corticosteroid) AND the patient has responded to Benlysta subcutaneous or intravenous. SLE-Initial-The patient has autoantibody-positive SLE, defined as positive for antinuclear antibodies [ANA] and/or anti-double-stranded DNA [anti-dsDNA] antibody AND Benlysta is being used concurrently with at least one other standard therapy (i.e., antimalarials [e.g., hydroxychloroquine], a systemic corticosteroid [e.g., prednisone], and/or other immunosuppressants [e.g., azathioprine, mycophenolate mofetil, methotrexate]) unless the patient is determined to be intolerant due to a significant toxicity,. Continuation-Benlysta is being used concurrently with at least one other standard therapy (i.e., antimalarials [e.g., hydroxychloroquine], a systemic corticosteroid [e.g., prednisone], and/or other immunosuppressants [e.g., azathioprine, mycophenolate mofetil, methotrexate]) unless the patient is determined to

PA Criteria	Criteria Details
	be intolerant due to a significant toxicity AND The patient has responded to Benlysta subcutaneous or intravenous.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BESREMI

Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other interferon products
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BETASERON/EXTAVIA

Products Affected

- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agent used for multiple sclerosis
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or after consultation with a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BEXAROTENE (ORAL)

Products Affected

- *bexarotene*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BEXAROTENE (TOPICAL)

Products Affected

- *bexarotene*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
Coverage Duration	1 year
Other Criteria	Adult T-Cell Leukemia/Lymphoma- approve if the patient has chronic/smoldering subtype and this medication is used as first-line therapy. Primary cutaneous B-Cell lymphoma-approve if used as skin-directed therapy for either (a or b): a) primary cutaneous marginal zone lymphoma or b) follicle center lymphoma.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Adult T-Cell Leukemia/Lymphoma, Primary Cutaneous B-Cell Lymphoma
Part B Prerequisite	No

BOSENTAN/AMBRISENTAN

Products Affected

- *ambrisentan*
- *bosentan oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) WHO Group 1, results of right heart cath
Age Restrictions	N/A
Prescriber Restrictions	For treatment of pulmonary arterial hypertension, ambrisentan or bosentan must be prescribed by or in consultation with a cardiologist or a pulmonologist. CTEPH - prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Authorization will be for 1 year.
Other Criteria	CTEPH - pt must have tried Adempas, has a contraindication to Adempas, or is currently receiving bosentan for CTEPH. Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chronic thromboembolic pulmonary hypertension (CTEPH) (bosentan)
Part B Prerequisite	No

BOSULIF

Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	ALL - 15 years and older. Myeloid/lymphoid neoplasms w eosinophilia- 18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	CML-approve if the patient has Ph-positive or BCR::ABL1-positive CML. For Ph-positive ALL-approve. Myeloid/lymphoid neoplasms with eosinophilia - approve if tumor has an ABL1 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Philadelphia chromosome positive Acute Lymphoblastic Leukemia, myeloid/lymphoid neoplasms with eosinophilia
Part B Prerequisite	No

BRAFTOVI

Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, BRAF V600 status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation. Colon or Rectal cancer- approve if the patient meets the following (A and B): A) The patient has BRAF V600E mutation-positive disease AND B) meets (i or ii): i) will be used as first-line systemic therapy for metastatic disease in combination with Erbitux (cetuximab intravenous infusion) and mFOLFOX6 (5-FU, leucovorin, and oxaliplatin) or ii) patient has previously received a chemotherapy regimen for colon or rectal cancer and this is prescribed as part of a combination regimen for colon or rectal cancer. NSCLC- approve if pt has BRAF V600E mutation-positive metastatic disease AND this medication will be taken in combination with Mektovi (binimatinib tablets). Appendiceal adenocarcinoma-approve if (A, B and C): A) BRAF V600E mutation-positive, and B) used as subsequent therapy for advanced or metastatic disease, and C) used in combination with Erbitux (cetuximab intravenous infusion) or Vectibix (panitumumab intravenous infusion).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Appendiceal adenocarcinoma
Part B Prerequisite	No

BRUKINSA

Products Affected

- BRUKINSA ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Follicular Lymphoma - approve if pt tried at least two other systemic regimens and will use this in combination with Gazyva (obinutuzumab intravenous infusion). Mantle Cell Lymphoma/CLL/SLL - approve. Marginal zone lymphoma-approve if the patient has tried at least one systemic regimen. Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma-approve. Hairy Cell Leukemia - approve if pt has received therapy for relapsed or refractory disease AND pt has progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Hairy Cell Leukemia
Part B Prerequisite	No

C1 ESTERASE INHIBITORS

Products Affected

- CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	prescribed by or in consultation with an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders
Coverage Duration	1 year
Other Criteria	Hereditary Angioedema (HAE) Due to C1 Inhibitor (C1-INH) Deficiency [Type I or Type II], Prophylaxis, Initial Therapy: approve if the patient has HAE type I or type II confirmed by low levels of functional C1-INH protein (less than 50 percent of normal) at baseline and lower than normal serum C4 levels at baseline. Patient is currently taking Cinryze for prophylaxis - approve if the patient meets the following criteria (i and ii): i) patient has a diagnosis of HAE type I or II, and ii) according to the prescriber, the patient has had a favorable clinical response since initiating Cinryze as prophylactic therapy compared with baseline.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CABLIVI

Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, concurrent medications
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	Approve for 12 months
Other Criteria	aTTP-approve if the requested medication was initiated in the inpatient setting in combination with plasma exchange therapy AND patient is currently receiving at least one immunosuppressive therapy AND if the patient has previously received Cablivi, he/she has not had more than two recurrences of aTTP while on Cablivi.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CABOMETYX

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, histology, RET gene rearrangement status for NSCLC
Age Restrictions	Neuroendocrine tumor/Thyroid carcinoma-12 years and older, other dx (except bone cancer)-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Renal Cell Carcinoma-Approve if the patient has relapsed or stage IV disease. Hepatocellular Carcinoma-approve if the patient has been previously treated with at least one other systemic therapy (e.g., Nexavar, Lenvima). Bone cancer-approve if the patient has Ewing sarcoma or osteosarcoma and has tried at least one previous systemic regimen. Thyroid carcinoma-approve if the patient has differentiated thyroid carcinoma, patient is refractory to radioactive iodine therapy and the patient has tried Lenvima or sorafenib. Endometrial carcinoma-approve if the patient has tried one systemic regimen. GIST-approve if the patient has tried two of the following-imatinib, Ayvakit, sunitinib, dasatinib, Stivarga or Qinlock. NSCLC-approve if the patient has RET rearrangement positive tumor. Neuroendocrine tumors- approve if (A and B): A) pt has locally advanced, unresectable, or metastatic disease, and B) meets (i or ii): i) patient has well-differentiated neuroendocrine tumors, or ii) patient has pancreatic or extra-pancreatic neuroendocrine tumors and the medication will be used as subsequent therapy. Adrenal gland tumor- approve if pt has locoregional unresectable or metastatic adrenocortical carcinoma. Pheochromocytoma/paraganglioma- approve if pt has locally unresectable disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Patients with Non-Small Cell Lung Cancer, Gastrointestinal stromal tumors (GIST), Bone cancer, Endometrial carcinoma, Adrenal gland tumor, Pheochromocytoma/paraganglioma
Part B Prerequisite	No

CALQUENCE

Products Affected

- CALQUENCE (ACALABRUTINIB
MAL)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	CLL and SLL-approve. Mantle Cell Lymphoma- approve if the patient meets (A or B): A) has tried at least one systemic regimen or is not a candidate for a systemic regimen (e.g., rituximab, dexamethasone, cytarabine, carboplatin, cisplatin, oxaliplatin, cyclophosphamide, doxorubicin, vincristine, prednisone, methotrexate, bendamustine, bortezomib, or lenalidomide) or B) this medication is used in combination with rituximab. Marginal Zone Lymphoma-approve if patient has tried at least one systemic regimen (e.g., bendamustine, rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone, lenalidomide, or chlorambucil). Waldenstrom Macroglobulinemia/Lymphoplasmacytic Lymphoma-approve if the patient has tried at least one systemic regimen (e.g., Brukinsa [zanubrutinib capsules], Imbruvica [ibrutinib tablets and capsules], rituximab, bendamustine, cyclophosphamide, dexamethasone, bortezomib, fludarabine, or cladribine)
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Waldenstrom's Macroglobulinemia/Lymphoplasmacytic Lymphoma, Marginal zone lymphoma.
Part B Prerequisite	No

CAMZYOS

Products Affected

- CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (initial and continuation)
Prescriber Restrictions	Prescribed by a cardiologist (initial and continuation)
Coverage Duration	Initial-8 months, continuation- 1 year
Other Criteria	Obstructive hypertrophic cardiomyopathy, initial-Approve if the pt meets the following criteria (i, ii and iii): i.Pt meets both of the following (a and b): a)Pt has at least 1 symptom associated w/obstructive hypertrophic cardiomyopathy (Note: examples include shortness of breath, chest pain, lightheadedness, fainting, fatigue, and reduced ability to perform physical exercise), AND b)Pt has New York Heart Association Class II or III symptoms of heart failure (Note:Class II signifies mild symptoms with moderate physical activity and some exercise limitations whereas Class III denotes noticeable symptoms with minimal physical activity and patients are only comfortable at rest), AND ii.Pt has left ventricular hypertrophy and meets 1 of the following (a or b): a)Pt has maximal left ventricular wall thickness greater than or equal to 15 mm, OR b)Pt has familial hypertrophic cardiomyopathy with a maximal left ventricular wall thickness greater than or equal to 13 mm, AND iii.Pt has a peak left ventricular outflow tract gradient greater than or equal to 50 mmHg (at rest or after provocation [Valsalva maneuver or post exercise]). Cont-Approve if pt meets ALL of the following criteria (i, ii, iii and iv): i.Pt has been established on therapy for at least 8 months (Note: pt who has received less than 8 months of therapy or who is restarting therapy is reviewed under initial therapy), AND ii.Pt meets both of the following (a and b): a)Currently or prior to starting therapy, pt has or has experienced at least 1 symptom associated with obstructive hypertrophic cardiomyopathy, AND

PA Criteria	Criteria Details
	<p>b)Currently or prior to starting therapy, pt is in or was in New York Heart Association Class II or III heart failure, AND iii.Pt has a current left ventricular ejection fraction of greater than or equal to 50 percent, AND iv.Pt meets at least 1 of the following (a or b): a)Pt experienced a beneficial clinical response when assessed by at least 1 objective measure (Note:Examples include improved peak oxygen consumption/mixed venous oxygen tension, decreases in left ventricular outflow tract gradient, reductions in N-terminal pro-B-type natriuretic peptide levels, decreased high-sensitivity cardiac troponin I levels, reduced ventricular mass index, and/or a reduction in maximum left atrial volume index), OR b)Pt experienced stabilization or improvement in at least 1 symptom related to obstructive hypertrophic cardiomyopathy (Note:Examples of symptoms include shortness of breath, chest pain, lightheadedness, fainting, fatigue, ability to perform physical exercise, and/or favorable changes in the Kansas City Cardiomyopathy Questionnaire-23 (KCCQ-23) Clinical Summary Score (CSS) or Hypertrophic Cardiomyopathy Symptom Questionnaire (HCMSQ) Shortness of Breath domain scores.)</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CAPRELSA

Products Affected

- CAPRELSA ORAL TABLET 100 MG,
300 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	MTC - approve. DTC - approve if refractory to radioactive iodine therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma.
Part B Prerequisite	No

CARGLUMIC ACID

Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a metabolic disease specialist or a specialist who focuses in the treatment of metabolic diseases
Coverage Duration	NAGS-Pt meets criteria no genetic test - 3mo. Pt had genetic test - 12mo, other-approve 7 days
Other Criteria	N-Acetylglutamate synthase deficiency with hyperammonemia-Approve if genetic testing confirmed a mutation leading to N-acetylglutamate synthase deficiency or if the patient has hyperammonemia. Propionic Acidemia or Methylmalonic Acidemia with Hyperammonemia, Acute Treatment-approve if the patient's plasma ammonia level is greater then or equal to 50 micromol/L and the requested medication will be used in conjunction with other ammonia-lowering therapies.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA) (generic carglumic acid)
Part B Prerequisite	No

CAYSTON

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist, infectious diseases specialist or a physician who specializes in the treatment of cystic fibrosis.
Coverage Duration	1 year
Other Criteria	Approve if the patient has <i>Pseudomonas aeruginosa</i> in culture of the airway (e.g., sputum culture, oropharyngeal culture, bronchoalveolar lavage culture).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CHEMET

Products Affected

- CHEMET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Blood lead level
Age Restrictions	Approve in patients between the age of 12 months and 18 years
Prescriber Restrictions	Prescribed by or in consultation with a professional experienced in the use of chelation therapy (eg, a medical toxicologist or a poison control center specialist)
Coverage Duration	Approve for 2 months
Other Criteria	Approve if Chemet is being used to treat acute lead poisoning (not as prophylaxis) and prior to starting Chemet therapy the patient's blood lead level was greater than 45 mcg/dL.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CIBINQO

Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a Biologic or with a Targeted Synthetic Disease-Modifying Antirheumatic Drug (DMARD). Concurrent use with an Anti-Interleukin Monoclonal Antibody. Concurrent use with other Janus Kinase Inhibitors. Concurrent use with a biologic immunomodulator. Concurrent use with other potent immunosuppressants.
Required Medical Information	Diagnosis
Age Restrictions	AD-12 years of age and older (initial therapy)
Prescriber Restrictions	Atopic Dermatitis-prescribed by or in consultation with an allergist, immunologist or dermatologist (initial therapy)
Coverage Duration	Initial-Atopic Dermatitis-3 months, Continuation-1 year
Other Criteria	Atopic Dermatitis, initial-approve if the patient has had a 4-month trial of at least one systemic therapy OR patient has tried at least one systemic therapy but was unable to tolerate a 4-month trial. Note: Examples of systemic therapies include Dupixent (dupilumab subcutaneous injection) or Adbry (tralokinumab-ldrm subcutaneous injection). Methotrexate, azathioprine, cyclosporine, and mycophenolate mofetil also count towards a trial of one systemic therapy. Continuation-Approve if the patient has been receiving Cibinvo for at least 90 days AND patient experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating Cibinvo) in at least one of the following: estimated body surface area affected, erythema, induration/papulation/edema, excoriations, lichenification, and/or a decreased requirement for other topical or systemic therapies for atopic dermatitis AND compared with baseline (prior to receiving Cibinvo), patient experienced an improvement in at least one symptom, such as decreased itching. Note: A patient who has received less than 3 months of therapy or who is restarting therapy with Cibinvo should be considered under initial therapy.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

CIMZIA

Products Affected

- CIMZIA POWDER FOR RECONST
- CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried
Age Restrictions	18 years and older for CD and PP (initial therapy). 2 years and older for JIA (initial therapy).
Prescriber Restrictions	All dx initial therapy only. RA, AS, JIA, prescribed by or in consultation with a rheumatologist. Crohn's disease, prescribed by or in consultation with a gastroenterologist. PsA prescribed by or in consultation with a rheumatologist or dermatologist. PP, prescribed by or in consultation with a dermatologist. nr-axSpA-prescribed by or in consultation with a rheumatologist
Coverage Duration	Approve through end of plan year
Other Criteria	AS initial tx, approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Xeljanz/XR, Cosentyx. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab product will also count. PsA initial tx, approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Cosentyx, Tremfya, a preferred ustekinumab product, Otezla, Orencia, Rinvoq, Skyrizi or Xeljanz/XR. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab/ustekinumab product will also count. RA initial tx, approve if the patient has tried two of the following drugs in the past: Enbrel, a preferred adalimumab product, Orencia, Rinvoq or Xeljanz/XR. Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab product will also count. CD initial tx, approve if patient has previously tried ONE of the following drugs in the past: a preferred adalimumab product, a preferred infliximab product, a preferred ustekinumab product, Skyrizi, Rinvoq, or Tremfya.

PA Criteria	Criteria Details
	<p>Note: if the patient does not meet this requirement, a previous trial of another non-preferred adalimumab/infliximab/ustekinumab product will also count. Plaque Psoriasis (PP), initial tx-approve if the patient has tried TWO of the following drugs in the past: Enbrel, a preferred adalimumab product, Skyrizi, a preferred ustekinumab product, Otezla, Cosentyx, Tremfya, Sotyktu. A trial of a non-preferred adalimumab/ustekinumab also counts. JIA initial tx, approve if the patient has tried TWO of the following drugs in the past: Enbrel, Orencia, Rinvoq/LQ, Xeljanz, a preferred adalimumab product. Note pt does not meet this requirement, a trial with a non-preferred adalimumab, Simponi Aria, tocilizumab, Kevzara, or infliximab will also count. Cont tx, AS/PsA/RA/CD/PP/JIA - approve if the pt had a response as determined by the prescriber. Non-radiographic axial spondylitis (nr-axSpA), initial tx-approve if the patient has objective signs of inflammation, defined as at least one of the following: C-reactive protein (CRP) elevated beyond the upper limit of normal for the reporting laboratory OR sacroilitis reported on MRI. nr-axSpA continuation tx-approve if the patient has had a response as determined by the prescriber. Please Note: preferred adalimumab products include Humira (NDCs starting with -00074), Cyltezo, Yuflyma. Preferred infliximab products include Remicade, Zymfentra. Preferred ustekinumab products include Stelara, Selarsdi, Yesintek.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CINACALCET

Products Affected

- *cinacalcet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Hypercalcemia d/t parathyroid CA-prescr/consult w/onco or endo. Hypercalcemia w/primary hyperparathyroidism-prescr/consult w/nephro or endo. Hyperparathyroidism in post-renal transplant-prescr/consult w/transplant physician/nephro/endo.
Coverage Duration	12 months
Other Criteria	Hypercalcemia due to parathyroid carcinoma-approve. Hypercalcemia in patients with primary hyperparathyroidism-approve if the patient has failed or is unable to undergo a parathyroidectomy due to a contraindication. Secondary Hyperparathyroidism in patients with chronic kidney disease on dialysis - deny under Medicare Part D (claim should be submitted under the ESRD bundles payment benefit). Hyperparathyroidism in Post-Renal Transplant Patients-approve if the baseline (prior to starting cinacalcet therapy) calcium and intact parathyroid hormone (iPTH) levels are above the normal range, as defined by the laboratory reference values.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	hyperparathyroidism in post-renal transplant patients
Part B Prerequisite	No

CLOBAZAM

Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*
- SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications tried
Age Restrictions	2 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with a neurologist (initial therapy)
Coverage Duration	1 year
Other Criteria	Lennox-Gastaut Syndrome, initial therapy-patient has tried and/or is concomitantly receiving one of the following: lamotrigine, topiramate, rufinamide, felbamate, Fintepla, Epidiolex or valproic acid. Treatment refractory seizures/epilepsy, initial therapy-patient has tried and/or is concomitantly receiving at least two other antiepileptic drugs (e.g., valproic acid, lamotrigine, topiramate, clonazepam, levetiracetam, zonisamide, felbamate). Continuation-prescriber confirms patient is responding to therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Dravet Syndrome and treatment-refractory seizures/epilepsy
Part B Prerequisite	No

COMETRIQ

Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)
- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis.
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	MTC - approve. Non-Small Cell Lung Cancer with RET Gene Rearrangements - approve. Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma-approve if the patient's carcinoma is refractory to radioactive iodine therapy and patient has tried a Vascular Endothelial Growth Factor Receptor (VEGFR)-targeted therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-Small Cell Lung Cancer with RET Gene Rearrangements, Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma
Part B Prerequisite	No

COPIKTRA

Products Affected

- COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Chronic Lymphocytic Leukemia/ Small Lymphocytic Lymphoma - approve if the patient has tried one systemic regimen (e.g., Imbruvica (ibrutinib capsules, tablets and oral solution), Venclexta (venetoclax tablets), rituximab, Gazyva (obinutuzumab intravenous infusion), chlorambucil, fludarabine, cyclophosphamide, bendamustine, high-dose methylprednisolone, Campath (alemtuzumab intravenous infusion), Calquence (acalabrutinib capsules), Brukinsa (zanubrutinib capsules), or Arzerra (ofatumumab intravenous infusion). T-cell lymphoma- For peripheral T-cell lymphoma, approve. For breast implant-associated anaplastic large cell lymphoma, or hepatosplenic T-cell lymphoma, approve if the patient has relapsed or refractory disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	T-cell Lymphoma
Part B Prerequisite	No

COSENTYX

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
Required Medical Information	Diagnosis and previous medications use
Age Restrictions	PP-6 yr and older. AS/Spondy/HS initial - 18 years of age and older. PsA-2 years and older. Enthesitis-4 years and older
Prescriber Restrictions	PP initial-presc/consult derm. PsA initial - prescribed by or in consultation with a dermatologist or rheumatologist. AS/spondylo/enthesitis initial- by or in consultation with rheumatologist. HS initial - by or in consult w/ dermatologist
Coverage Duration	Approve through end of plan year
Other Criteria	<p>INITIAL THERAPY: HIDRADENITIS SUPPURATIVA (HS): tried at least one other therapy (e.g. systemic antibiotics, isotretinoin). NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS: objective signs of inflammation and meets a or b: a) C-reactive protein elevated beyond the upper limit of normal or b) sacroiliitis reported on MRI. PLQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent (e.g., methotrexate [MTX], cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial of at least one biologic that is not Cosentyx or a Cosentyx biosimilar also counts) or B) contraindication to MTX. CONTINUATION THERAPY: ALL INDICATIONS: patient has experienced benefit from the medication.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

COTELLIC

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Melanoma (unresectable, advanced or metastatic) - being prescribed in combination with Zelboraf AND patient has BRAF V600 mutation positive disease. CNS Cancer-approve if the patient has BRAF V600 mutation-positive disease AND medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma/neuroglioma/glioneuronal tumor OR ii. Recurrent or progressive disease for one of the following conditions (a or b): a) glioma/circumscribed glioma OR b) Glioblastoma, OR iii. Melanoma with brain metastases AND medication with be taken in combination with Zelboraf (vemurafenib tablets). Histiocytic Neoplasm-approve if the patient meets one of the following (i, ii, or iii): i. Patient has Langerhans cell histiocytosis and one of the following (a, b, or c): a) Multisystem disease OR b) Pulmonary disease OR c) Central nervous system lesions OR ii. Patient has Erdheim Chester disease OR iii. Patient has Rosai-Dorfman disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Central Nervous System Cancer
Part B Prerequisite	No

CRESEMBA (ORAL)

Products Affected

- CRESEMBA ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Candidiasis of the esophagus - HIV infection, sepsis
Part B Prerequisite	No

CYSTEAMINE (OPHTHALMIC)

Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an ophthalmologist or a metabolic disease specialist or specialist who focuses in the treatment of metabolic diseases
Coverage Duration	1 year
Other Criteria	Approve if the patient has corneal cysteine crystal deposits confirmed by slit-lamp examination
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CYSTEAMINE (ORAL)

Products Affected

- CYSTAGON

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Cystagon and Procysbi
Required Medical Information	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist or a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
Coverage Duration	1 year
Other Criteria	Cystinosis, nephropathic-approve if the prescriber confirms the diagnosis was confirmed by genetic testing confirming a mutation of the CTNS gene OR white blood cell cystine concentration above the upper limit of the normal reference range for the reporting laboratory.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DALFAMPRIDINE

Products Affected

- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older (initial and continuation therapy)
Prescriber Restrictions	MS. If prescribed by, or in consultation with, a neurologist or MS specialist (initial and continuation).
Coverage Duration	Initial-4months, Continuation-1 year
Other Criteria	Initial-approve if the patient is ambulatory, the requested medication is being used to improve or maintain mobility in a patient with MS and the patient has impaired ambulation as evaluated by an objective measure (e.g., timed 25 foot walk and multiple sclerosis walking scale-12). Continuation-approve if the patient is ambulatory, the requested medication is being used to improve or maintain mobility in a patient with MS and the patient has responded to or is benefiting from therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DANZITEN

Products Affected

- DANZITEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	ALL - 15 years and older, GIST/Myeloid/lymphoid neoplasms/melanoma, cutaneous-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	CML- approve if BCR::ABL1-mutation positive or Philadelphia chromosome positive. ALL-Philadelphia chromosome positive. Pigmented villonodular synovitis/tenosynovial giant cell tumor-patient has tried Turalio or Romvimza or cannot take Turalio or Romvimza. For GIST, approve if the patient has tried two of the following: imatinib, avapritinib, sunitinib, dasatinib, regorafenib or ripretinib. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an ABL1 rearrangement. For melanoma, cutaneous - approve if the patient has metastatic or unresectable disease AND has an activating KIT mutation AND has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Acute lymphoblastic leukemia, Pigmented villonodular synovitis/Tenosynovial giant cell tumor, GIST, cutaneous melanoma and myeloid/lymphoid neoplasms with eosinophilia
Part B Prerequisite	No

DAURISMO

Products Affected

- DAURISMO ORAL TABLET 100 MG,
25 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medications that will be used in combination, comorbidities
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML - approve if Daurismo will be used in combination with cytarabine.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DEFERASIROX

Products Affected

- *deferasirox*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Serum ferritin level
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	1 year
Other Criteria	Transfusion-related chronic iron overload, initial therapy - approve if the patient is receiving blood transfusions at regular intervals for various conditions (eg, thalassemia syndromes, myelodysplastic syndrome, chronic anemia, sickle cell disease) AND prior to starting therapy, the serum ferritin level is greater than 1,000 mcg/L. Non-transfusion-dependent thalassemia syndromes chronic iron overload, initial therapy - approve if prior to starting therapy the serum ferritin level is greater than 300 mcg/L. Continuation therapy - approve if the patient is benefiting from therapy as confirmed by the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DEFERIPRONE

Products Affected

- *deferiprone*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Serum ferritin level
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	1 year
Other Criteria	Iron overload, chronic-transfusion related due to thalassemia syndrome or related to sickle cell disease or other anemias Initial therapy - approve. Continuation therapy - approve is the patient is benefiting from therapy as confirmed by the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIABETIC SUPPLY - ALCOHOL PADS

Products Affected

- *alcohol pads*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIABETIC SUPPLY - GAUZE PADS

Products Affected

- GAUZE PADS 2 X 2

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIABETIC SUPPLY - NEEDLES

Products Affected

- NOVO PEN NEEDLE
- EMBECTA PEN NEEDLE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIABETIC SUPPLY - SYRINGES

Products Affected

- EMBECTA INSULIN SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Approve if the prescriber confirms that the medical supply is being requested for a use that is directly associated with delivering insulin to the body.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIACOMIT

Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	6 months and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with an neurologist (initial therapy)
Coverage Duration	1 year
Other Criteria	Dravet Syndrome-Initial therapy-approve if the patient is concomitantly receiving clobazam or is unable to take clobazam due to adverse events. Dravet Syndrome-Continuation-approve if the patient is responding to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIMETHYL FUMARATE

Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS).
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DOPTELET

Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Thrombocytopenia in chronic liver disease - 18 years and older
Prescriber Restrictions	Chronic ITP-prescribed by or after consultation with a hematologist (initial therapy)
Coverage Duration	Thrombo w/chronic liver disease-5 days, chronic ITP initial-3 months, cont-1 year
Other Criteria	THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE (A and B): A) current platelet count less than 50 x 10 ⁹ /L and B) scheduled to undergo a procedure within 10 to 13 days after starting Doptelet therapy. CHRONIC ITP, INITIAL THERAPY (A and B): A): (i or ii): i) platelet count less than 30,000 microliters or ii) platelet count less than 50,000 microliters and patient is at an increased risk of bleeding, and B) tried one other therapy (e.g., systemic corticosteroids, intravenous immunoglobulin, anti-D immunoglobulin, eltrombopag tablets and oral suspension, romiplostim subcutaneous injection, fostamatinib tablets, rituximab) or had a splenectomy. CHRONIC ITP, CONTINUATION THERAPY: patient had beneficial clinical response and remains at risk for bleeding complications.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DROXIDOPA

Products Affected

- *droxidopa*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Medication history (as described in Other Criteria field)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a neurologist
Coverage Duration	12 months
Other Criteria	NOH, approve if the patient meets ALL of the following criteria: a) Patient has been diagnosed with symptomatic NOH due to primary autonomic failure (Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy, AND b) Patient has tried midodrine
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DUAL OREXIN RECEPTOR ANTAGONIST

Products Affected

- BELSOMRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Treatment of insomnia, characterized by difficulties with sleep onset and /or sleep maintenance-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DUPIXENT

Products Affected

- DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML
- DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Xolair or another Anti-interleukin (IL) Monoclonal Antibody (i.e., Adbry, Cinqair, Fasenra, Nucala, Tezspire, or Xolair). Concurrent use with Janus Kinase Inhibitors (JAKis) [oral or topical].
Required Medical Information	Diagnosis, prescriber specialty, other medications tried and length of trials. COPD INITIAL: meets (all of A, B, and C): A) blood eosinophil at least 300 cells per microliter within previous 6 weeks or prior to Dupixent or another monoclonal antibody, and B) received at least 3 months of combination therapy with at least two of LAMA, LABA or ICS, and, and C) meets (i or ii): i) two or more COPD exacerbations in previous 12 months requiring systemic CS or antibiotics or ii) COPD exacerbation requiring hospitalization in previous 12 months. COPD CONTINUATION (all of A, B and C): A) received Dupixent for at least 6 months and B) continues LABA and LAMA, and C) beneficial response (e.g. reduced symptoms, exacerbations, hospitalizations, ED/urgent care visits, improved lung function). CSU INITIAL: urticaria for greater than 6 weeks (prior to Dupixent), with symptoms at least 3 days/week despite daily non-sedating H1 antihistamine tx. CSU CONTINUATION (A and B): A) received at least 6 months of Dupixent and B) experienced a beneficial clinical response, defined by decreased itch severity, decreased number of hives or decreased size of hives. Bullous Pemphigoid Initial: Approve. Bullous Pemphigoid Continuous: meets both (i and ii): i. received at least 6 months of therapy with Dupixent AND ii. experienced a beneficial clinical response, defined by decreased area of skin involvement, lesions, including blisters or erosions (bullae), urticaria, erythema, or reduced or no need for systemic or topical corticosteroid therapy.
Age Restrictions	Initial therapy only: AD-6 months and older, asthma-6 years of age and older, Esophagitis-1 yr and older, Chronic Rhinosinusitis/CSU- 12 and older, Prurigo nodularis/COPD/BP-18 and older
Prescriber Restrictions	Initial therapy only: Atopic Dermatitis/prurigo nodularis/CSU-Prescribed by or in consultation with an allergist, immunologist or dermatologist, asthma-prescribed by or in consultation with an allergist, immunologist or pulmonologist. Rhinosinusitis-prescribed by or in consultation with an

PA Criteria	Criteria Details
	allergist, immunologist or otolaryngologist. Esophagitis-presc/consult-allergist or gastro. COPD-prescribed by or in consultation with an allergist, immunologist, or pulmonologist. BP-prescr/consult with derm
Coverage Duration	AD-Init-4mo, Cont-1 yr, asthma/Rhinosinusitis/esophagitis/PN/COPD/CSU/BP-init-6 mo, cont 1 yr
Other Criteria	<p>INITIAL CRITERIA: AD: tried at least 1 medium to super-high-potency topical corticosteroid (CS), unless topical CS therapy not advisable or pt is less than 2 years old. ASTHMA (all of A, B, and C): A) blood eosinophil greater than or equal to 150 cells per microliter within previous 6 weeks or within 6 weeks prior to Dupixent or another monoclonal antibody or has oral CS-dependent asthma, B) used an ICS in combination with at least one additional asthma controller/maintenance medication, and C) uncontrolled asthma prior to any asthma monoclonal antibody as defined by one of the following (one of a, b, c, d, or e): a) two or more asthma exacerbations requiring oral CS in the past year, b) one or more asthma exacerbations requiring hospital/urgent care/ED visit in the past year, c) FEV1 less than 80 percent predicted (adults only), d) FEV1/FVC less than 0.8, or e) worsened asthma with oral CS taper. CRSwNP (all of A, B, C and D): A) concurrent use with nasal CS, B) presence of at least two of the following symptoms for 6 months: nasal congestion, nasal obstruction, nasal discharge, reduction/loss of smell, C) received oral CS at least 5 days in last 2 years (unless contraindicated) or patient had prior surgery for nasal polyps, and D) diagnosis confirmed by direct exam, endoscopy, or sinus CT. EoE (all of A, B, C, and D): A) weighs 15 kg or more, B) endoscopic biopsy demonstrating greater than or equal to 15 intraepithelial eosinophils per high-power field, C) does not have a secondary cause of EoE, and D) received an Rx-strength PPI for at least 8 weeks. PRURIGO NODULARIS: approve if pt has pruritus lasting at least 6 weeks.</p> <p>CONTINUATION CRITERIA: AD: responding positively to therapy. ASTHMA: responding positively to therapy and concurrent use with ICS. CRSwNP (all of A, B, and C): A) received Dupixent for at least 6 months, B) responding positively to therapy, and C) concurrent use with intranasal CS. EoE (A and B): A) received Dupixent for at least 6 months and B) reduction in intraepithelial eosinophil count, decreased dysphagia/pain upon swallowing, or reduced frequency/severity of food impaction. PRURIGO NODULARIS (A and B): A) received Dupixent for at least 6 months and B) reduction in nodular lesion count, pruritis, or nodular lesion size.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

EMGALITY

Products Affected

- EMGALITY PEN
- EMGALITY SUBCUTANEOUS SYRINGE 120 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	Combination therapy with another cGRP inhibitor for migraine headache prevention
Required Medical Information	Diagnosis, number of migraine or cluster headaches per month
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Migraine headache prevention-Approve if the patient meets the following criteria (A and B): A) Patient has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication), AND B) If pt is currently taking Emgality, pt has had significant clinical benefit from the medication. Examples of significant clinical benefit include a reduction in the overall number of migraine days per month or a reduction in number of severe migraine days per month from the time that Emgality was initiated. Episodic cluster headache treatment-approve if the patient has between one headache every other day and eight headaches per day.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with biologic therapy or targeted synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried.
Age Restrictions	PP-4 years and older (initial therapy)
Prescriber Restrictions	Initial only-RA/AS/JIA/JRA,prescribed by or in consult w/ rheumatologist. Psoriatic arthritis, prescribed by or in consultation w/ rheumatologist or dermatologist.Plaque psoriasis (PP), prescribed by or in consult w/ dermatologist.GVHD,prescribed by or in consult w/ oncologist,hematologist,or physician affiliated w/ transplant center.Behcet's disease,prescribed by or in consult w/ rheumatologist,dermatologist,ophthalmologist,gastroenterologist,or neurologist.
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: RHEUMATOID ARTHRITIS (RA): tried one conventional synthetic DMARD for at least 3 months (see Note 1). JUVENILE IDIOPATHIC ARTHRITIS (JIA)/JRA (one of A, B, C, or D): A) patient has aggressive disease, B) tried one other systemic therapy (e.g., methotrexate [MTX], sulfasalazine, leflunomide, NSAID, or a biologic that is not a biosimilar of the requested product), C) patient will be started on Enbrel concurrently with MTX, sulfasalazine, or leflunomide, or D) patient has an absolute contraindication to MTX, sulfasalazine, or leflunomide. PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent for at least 3 months, unless intolerant (e.g., MTX, cyclosporine, Soriatane, oral methoxsalen plus PUVA) [see Note 1] or B) patient has a contraindication to one oral agent for psoriasis such as MTX. GRAFT VERSUS HOST DISEASE (GVHD): approve. BEHCET'S: tried at least one conventional therapy (e.g., systemic corticosteroid, immunosuppressant, interferon alfa, mycophenolate), adalimumab, or

PA Criteria	Criteria Details
	infliximab. CONTINUATION THERAPY: ALL INDICATIONS: patient had a response to therapy. Note 1: a biologic that is not a biosimilar of the requested product will also count.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Graft versus host disease (GVHD), Behcet's disease
Part B Prerequisite	No

ENDARI

Products Affected

- *glutamine (sickle cell)*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prescriber specialty
Age Restrictions	Greater than or equal to 5 years of age
Prescriber Restrictions	Prescribed by, or in consultation with, a physician who specializes in sickle cell disease (e.g., a hematologist)
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EPIDIOLEX

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies
Age Restrictions	Patients 1 year and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with a neurologist (initial therapy)
Coverage Duration	1 year
Other Criteria	Dravet Syndrome-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs or if the patient has tried or is concomitantly receiving one of Diacomit or clobazam or Fintepla. Lennox Gastaut Syndrome-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs. Tuberous Sclerosis Complex-approve if the patient has tried or is concomitantly receiving at least two other antiseizure drugs. Continuation of therapy-approve if the patient is responding to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EPOETIN ALFA

Products Affected

- PROCRIT INJECTION SOLUTION
10,000 UNIT/ML, 2,000 UNIT/ML,
20,000 UNIT/ML, 3,000 UNIT/ML, 4,000
UNIT/ML, 40,000 UNIT/ML
- RETACRIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	MDS anemia = 18 years of age and older
Prescriber Restrictions	MDS anemia, myelofibrosis- prescribed by or in consultation with, a hematologist or oncologist.
Coverage Duration	Chemo-6m, Transfus-1m, CKD-1yr, Myelofibrosis-init-3 mo, cont-1 yr, all others-1 yr
Other Criteria	Anemia in a pt with Chronic Kidney Disease (CKD) not on dialysis- for initial therapy, approve if hemoglobin (Hb) is less than 10.0 g/dL for adults or less than or equal to 11 g/dL for children, or for continuation of therapy in a pt currently on an erythropoiesis-stimulating agent (ESA) approve if Hb is less than or equal to 12 g/dL. Anemia in a pt with cancer due to chemotherapy- approve if pt is currently receiving myelosuppressive chemo as a non-curative treatment and (for initial therapy) Hb is less than 10.0 g/dL or (if currently on ESA) Hb is less than or equal to 12.0 g/dL. Anemia in HIV with zidovudine- for initial therapy, approve if Hb is less than 10.0 g/dL or serum erythropoietin level is 500 mU/mL or less, or for continuation of therapy in a pt currently on ESA, approve if Hb is 12.0 g/dL or less. Surgical pts to reduce RBC transfusions - Approve if Hb is less than or equal to 13, AND surgery is elective, nonvascular and non-cardiac AND pt is unwilling or unable to donate autologous blood prior to surgery. MDS- for initial therapy, approve if Hb is less than 10 g/dL or serum erythropoietin level is 500 mU/mL or less, or for continuation of therapy in a pt currently on ESA approve if Hb is 12.0 g/dL or less. Myelofibrosis- for Initial therapy approve if patient has a Hb less than 10 or serum erythropoietin less than or equal to 500 mU/mL, or for continuation of therapy in pt currently on ESA hemoglobin is less than or

PA Criteria	Criteria Details
	equal to 12g/dL. Anemia in patients with chronic renal failure on dialysis - deny under Medicare Part D (claim should be submitted under the ESRD bundled payment benefit).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Anemia due to myelodysplastic syndrome (MDS), Myelofibrosis
Part B Prerequisite	No

ERIVEDGE

Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	BCC (La or Met) - must not have had disease progression while on Odomzo.
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Basal cell carcinoma, locally advanced-approve. Central nervous system cancer (this includes brain and spinal cord tumors)-approve if the patient has medulloblastoma, the patient has tried at least one chemotherapy agent and according to the prescriber, the patient has a mutation of the sonic hedgehog pathway. Basal cell carcinoma, metastatic (this includes primary or recurrent nodal metastases and distant metastases)-approve. Diffuse Basal Cell Carcinoma Formation, including basal cell nevus syndrome (Gorlin syndrome) or other genetic forms of multiple basal cell carcinoma - approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Central nervous System Cancer, diffuse basal cell carcinoma formation
Part B Prerequisite	No

ERLEADA

Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate cancer-non-metastatic, castration resistant and prostate cancer-metastatic, castration sensitive-approve if the requested medication will be used in combination with a gonadotropin-releasing hormone (GnRH) analog [for example: leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets)] or if the patient has had a bilateral orchiectomy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ERLOTINIB

Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Advanced or Metastatic NSCLC, approve if the patient has EGFR mutation positive non-small cell lung cancer as detected by an approved test. Note- Examples of EGFR mutation-positive non-small cell lung cancer include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S768I. RCC, approve if the patient has recurrent or advanced non-clear cell histology RCC or if the patient had hereditary leiomyomatosis and renal cell carcinoma and erlotinib will be used in combination with bevacizumab. Bone cancer-approve if the patient has chordoma and has tried at least one previous therapy. Pancreatic cancer-approve if the medication is used in combination with gemcitabine and if the patient has locally advanced, metastatic or recurrent disease. Vulvar cancer-approve if the patient has advanced, recurrent or metastatic disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Renal Cell Carcinoma, vulvar cancer and Bone Cancer-Chordoma.
Part B Prerequisite	No

EVEROLIMUS

Products Affected

- *everolimus (antineoplastic) oral tablet*
- *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*
- *torpenz*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer-HER2 status, hormone receptor (HR) status.
Age Restrictions	All dx except TSC associated SEGA, renal angiomyolipoma or partial onset seizures-18 years and older.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Breast Cancer-approve if pt meets ALL the following (A, B, C, D, E, and F):A)pt has recurrent or metastatic,HR+ disease AND B)pt has HER2-negative breast cancer AND C)pt has tried at least 1 prior endocrine therapy AND D)pt meets 1 of the following conditions (i or ii):i.pt is a postmenopausal woman or man OR ii.pt is pre/perimenopausal woman AND is receiving ovarian suppression/ablation with a GnRH agonist, or has had surgical bilateral oophorectomy or ovarian irradiation AND E)pt meets 1 of the following conditions (i or ii): i.Everolimus will be used in combo with exemestane and pt meets 1 of the following:pt is male and is receiving a GnRH analog or pt is a woman or ii. Everolimus will be used in combo with fulvestrant or tamoxifen AND F)pt has not had disease progression while on Everolimus. RCC, relapsed or Stage IV disease-approve if using for non-clear cell disease or if using for clear cell disease, pt has tried 1 prior systemic therapy (e.g., Inlyta, Votrient, Sutent, Cabometyx, Nexavar).TSC Associated SEGA-approve if pt requires therapeutic intervention but cannot be curatively resected. Thymomas and Thymic Carcinomas-approve if pt has tried chemotherapy or cannot tolerate chemotherapy.TSC associated renal angiomyolipoma -approve. WM/LPL - approve if pt has progressive or relapsed disease or if pt has not responded to primary therapy. Thyroid Carcinoma, differentiated-approve if pt is refractory to radioactive iodine therapy. Endometrial Carcinoma-

PA Criteria	Criteria Details
	<p>approve if everolimus will be used in combo with letrozole. GIST-approve if pt has tried 2 of the following drugs: Sutent, Sprycel, Stivarga, Ayvakit, Qinlock or imatinib AND there is confirmation that everolimus will be used in combo with 1 of these drugs (Sutent, Stivarga, or imatinib) in the treatment of GIST. TSC-associated partial-onset seizures-approve. NET tumors of the pancreas, GI Tract, Lung and Thymus (carcinoid tumors)-approve. Soft tissue sarcoma-approve if pt has perivascular epithelioid cell tumors (PE Coma) or recurrent angiomyolipoma/lymphangioleiomyomatosis. Classic hodgkin lymphoma-approve if pt has relapsed or refractory disease AND has tried at least three prior lines of chemotherapy. Histiocytic neoplasm-approve if pt has Erdheim-Chester disease or, Rosai-Dorfman disease or Langerhans cell histiocytosis. Patient must also have PIK3CA mutation. Meningioma-approve if pt has recurrent or progressive disease AND pt has surgically inaccessible disease and radiation therapy is not possible AND medication will be used in combination with a somatostatin analogue. Uterine Sarcoma-approve if the patient has advanced, recurrent, metastatic, or inoperable disease, AND has a perivascular epithelioid cell tumor (PEComa), AND has tried at least one systemic regimen. Note: Examples of systemic regimen include doxorubicin, docetaxel, gemcitabine, ifosfamide, dacarbazine.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	neuroendocrine tumors of the thymus (Carcinoid tumors). Soft tissue sarcoma, classical Hodgkin lymphoma, Waldenstrom's Macroglobulinemia/Lymphoplasmacytic Lymphoma (WM/LPL), Thymomas and Thymic carcinomas, Differentiated Thyroid Carcinoma, Endometrial Carcinoma, Gastrointestinal Stromal Tumors (GIST), men with breast cancer, Pre-peri-menopausal women with breast cancer, Histiocytic Neoplasm, uterine sarcoma, meningioma
Part B Prerequisite	No

FASENRA

Products Affected

- FASENRA PEN
- FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another monoclonal antibody therapy.
Required Medical Information	Diagnosis
Age Restrictions	Asthma: 6 years of age and older, EGPA: 18 years and older
Prescriber Restrictions	Asthma: Prescribed by or in consultation with an allergist, immunologist, or pulmonologist. EGPA: Prescribed by or in consultation with an allergist, immunologist, pulmonologist, or rheumatologist
Coverage Duration	Asthma: 6 months initial, 12 months continuation. EGPA: 8 months initial, 12 months continuation.
Other Criteria	<p>INITIAL THERAPY: ASTHMA (all of A, B, and C): A) blood eosinophil greater than or equal to 150 cells per microliter within previous 6 weeks or within 6 weeks prior to Fasenra or another monoclonal antibody, B) used an inhaled corticosteroid (ICS) in combination with at least one additional asthma controller/maintenance medication, and C) uncontrolled asthma prior to any asthma monoclonal antibody as defined by one of the following (a, b, c, d, or e): a) one or more exacerbations requiring a systemic CS in the past year, b) one or more exacerbations requiring hospital/urgent care/emergency department visit in the past year, c) FEV1 less than 80 percent predicted or less than 90 percent predicted for patients less than 18, d) FEV1/FVC less than 0.80, or e) worsened asthma with systemic CS taper. EGPA: (all of A, B, and C): A) active disease, and B) currently on systemic CS for at least 4 weeks, and C) blood eosinophil greater than or equal to 150 cells per microliter within previous 4 weeks or prior to treatment with any monoclonal antibody that may alter eosinophil levels.</p> <p>CONTINUATION THERAPY: ASTHMA (A and B): A) patient has responded to therapy (e.g., decrease in any of the following: asthma exacerbations, asthma symptoms, hospitalizations, emergency department/urgent care visits, physician visits, requirement for oral corticosteroid therapy) and B) continues to receive therapy with an ICS.</p>

PA Criteria	Criteria Details
	EGPA: patient has responded to therapy (e.g. reduced rate of relapse, CS dose reduction, reduced eosinophil levels).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FINGOLIMOD

Products Affected

- *fingolimod*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of fingolimod with other disease-modifying agents used for multiple sclerosis (MS).
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	10 years and older
Prescriber Restrictions	Prescribed by, or in consultation with, a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FINTEPLA

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	2 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with an neurologist (initial therapy)
Coverage Duration	1 year
Other Criteria	Dravet Syndrome-Initial therapy-approve if the patient has tried or is concomitantly receiving at least two other antiepileptic drugs or patient has tried or is concomitantly receiving Epidiolex, Clobazam or Diacomit. Dravet Syndrome-Continuation-approve if the patient is responding to therapy. Lennox-Gastaut Syndrome, initial-approve if the patient has tried or is concomitantly receiving at least two other antiepileptic drugs. Lennox-Gastaut Syndrome, continuation-approve if the patient is responding to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FIRMAGON

Products Affected

- FIRMAGON KIT W DILUENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a oncologist
Coverage Duration	1 year
Other Criteria	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Patients new to therapy, are required to try Eligard or Orgovyx prior to approval of Firmagon.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FOTIVDA

Products Affected

- FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Renal Cell Carcinoma (RCC)-approve if the patient has relapsed or Stage IV disease and has tried at least two other systemic regimens. Note: Examples of systemic regimens for renal cell carcinoma include axitinib tablets, axitinib + pembrolizumab injection, cabozantinib tablets, cabozantinib + nivolumab injection, sunitinib malate capsules, pazopanib tablets, sorafenib tablets, and lenvatinib capsules + everolimus.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FRUZAQLA

Products Affected

- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Colon cancer, rectal cancer, or appendiceal cancer-Approve if the patient meets the following (A and B): A.Patient has advanced or metastatic disease, AND B.Patient has previously been treated with the following (i, ii, and iii): i.Fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, Note: Examples of fluoropyrimidine agents include 5-fluorouracil (5-FU) and capecitabine. AND ii.An anti-vascular endothelial growth factor (VEGF) agent, Note: Examples of anti-VEGF agents include bevacizumab. AND iii. If the tumor is RAS wild-type (KRAS wild-type and NRAS wild-type) [that is, the tumor or metastases are KRAS and NRAS mutation negative], the patient meets ONE of the following (a or b): a.According to the prescriber, anti-epidermal growth factor receptor (EGFR) therapy is NOT medically appropriate, OR b. The patient has received an anti-EGFR therapy. Note: Examples of anti-EGFR therapy includes Erbitux (cetuximab intravenous infusion) and Vectibix (panitumumab intravenous infusion).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Appendiceal cancer
Part B Prerequisite	No

FULPHILA

Products Affected

- FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Cancer patients receiving chemotherapy-prescribed by or in consultation with an oncologist or hematologist. PBPC - prescribed by or in consultation with an oncologist, hematologist, or physician that specializes in transplantation.
Coverage Duration	Cancer pts receiving chemo-6 mo. PBPC-30 days.
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient meets one of the following: 1) is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), 2) patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia according to the prescribing physician (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status or HIV infection, or 3) patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor and a reduced dose or frequency of chemotherapy may compromise treatment.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients undergoing PBPC collection and therapy
Part B Prerequisite	No

GATTEX

Products Affected

- GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	1 year and older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist (initial and continuation)
Coverage Duration	1 year
Other Criteria	Initial-approve if the patient is currently receiving parenteral nutrition on 3 or more days per week or according to the prescriber, the patient is unable to receive adequate total parenteral nutrition required for caloric needs. Continuation-approve if the patient has experienced improvement.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GAVRETO

Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	NSCLC-18 years and older, thyroid cancer-12 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NSCLC-approve if the patient has advanced, recurrent, or metastatic disease and rearranged during transfection (RET) fusion-positive disease detected by an Food and Drug Administration (FDA) approved test. Differentiated Thyroid Cancer- pt has unresectable, recurrent, or metastatic disease AND pt has RET fusion-positive or RET-mutation-positive disease AND disease requires treatment with systemic therapy AND the disease is radioactive iodine-refractory. Anaplastic thyroid cancer or Medullary Thyroid Cancer- pt has unresectable, recurrent, or metastatic disease AND pt has RET fusion-positive or RET-mutation-positive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Medullary Thyroid Cancer, Anaplastic Thyroid Cancer
Part B Prerequisite	No

GEFITINIB

Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NSCLC-approve if the patient has advanced or metastatic disease and the patient has sensitizing EGFR mutation-positive NSCLC as detected by an approved test. Note: Examples of sensitizing EGFR mutation-positive NSCLC include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S768I.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	NSCLC with EGFR L861Q, G719X, or S768I mutations.
Part B Prerequisite	No

GIOTRIF

Products Affected

- GIOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For NSCLC - EGFR exon deletions or mutations, or if NSCLC is squamous cell type
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	NSCLC EGFR pos - For the treatment of advanced or metastatic non small cell lung cancer (NSCLC)-approve if the patient has sensitizing EGFR mutation-positive NSCLC as detected by an approved test. Note: examples of sensitizing EGFR mutation-positive NSCLC include the following mutations : exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S768I. NSCLC metastatic squamous cell must have disease progression after treatment with platinum based chemotherapy. Head and neck cancer-approve if the patient has non-nasopharyngeal head and neck cancer and the patient has disease progression on or after platinum based chemotherapy. (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan)
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Head and neck cancer
Part B Prerequisite	Yes

GLATIRAMER

Products Affected

- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agent used for multiple sclerosis
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or after consultation with a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GLUCAGON-LIKE PEPTIDE-1 AGONISTS

Products Affected

- *exenatide subcutaneous pen injector 10 mcg/dose(250 mcg/ml) 2.4 ml, 5 mcg/dose (250 mcg/ml) 1.2 ml* MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- MOUNJARO
- OZEMPI SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 RYBELSUS
- TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GOMEKLI

Products Affected

- GOMEKLI ORAL CAPSULE 1 MG, 2 MG
- GOMEKLI ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	2 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NEUROFIBROMATOSIS TYPE 1- patient has or had symptomatic plexiform neurofibromas prior to starting Gomekli and the tumor is not amenable to complete resection.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GONADOTROPIN-RELEASING HORMONE AGONISTS - ONCOLOGY

Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)
- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prostate cancer- prescribed by or in consultation with an oncologist or urologist. Head and neck-salivary gland tumors- prescribed by or in consultation with an oncologist.
Coverage Duration	1 year
Other Criteria	Head and neck cancer-salivary gland tumor- approve if pt has recurrent, unresectable, or metastatic disease AND androgen receptor-positive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Head and neck cancer- salivary gland tumors (Eligard only)
Part B Prerequisite	No

GRALISE/HORIZANT/LYRICA CR

Products Affected

- *gabapentin oral tablet extended release 24 hr 300 mg, 600 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GROWTH HORMONES

Products Affected

- OMNITROPE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	GHD in Children/Adolescents. Pt meets one of the following-1-had 2 GH stim tests with the following-levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon and both are less than 10ng/mL OR had at least 1 GH test and results are less than 10ng/mL and has at least one risk factor for GHD (e.g., ht for age curve deviated down across 2 major height percentiles [e.g., from above the 25 percentile to below the 10 percentile], growth rate is less than the expected normal growth rate based on age and gender, low IGF-1 and/or IGFBP-3 levels). 2. brain radiation or tumor resection and pt has 1 GH stim test less than 10ng/mL or has def in at least 1 other pituitary hormone (that is, ACTH, TSH, gonadotropin deficiency [LH and/or FSH] are counted as 1 def], or prolactin).3. congenital hypopituitarism and has one GH stim test less than 10ng/mL OR def in at least one other pituitary hormone and/or the patient has the imaging triad of ectopic posterior pituitary and pituitary hypoplasia with abnormal pituitary stalk 4.pt has multiple pituitary deficiencies and pt has 3 or more pituitary hormone deficiencies or pt has had one GH test less than 10ng/mL 5.pt had a hypophysectomy. Cont-pt responding to therapy
Age Restrictions	ISS 5 y/o or older, SGA 2 y/o or older, SBS 18 y/o or older
Prescriber Restrictions	GHD (Initial tx children or adolescents w/o hypophysectomy), GHD adults or transitional adolescents, Noonan (initial), Prader Willi (initial for child/adult and cont tx in adults), SHOX (initial), SGA (initial) - prescribed by or in consultation with an endocrinologist. CKD (initial) endocrinologist or nephrologist.
Coverage Duration	ISS - 6 mos initial, 12 months cont tx, SBS 1 month, others 12 mos
Other Criteria	GHD initial in adults and adolescents 1. endocrine must certify not being prescribed for anti-aging or to enhance athletic performance, 2. has either childhood onset or adult onset resulting from GHD alone, multiple hormone deficiency from pituitary dx, hypothalamic dz, pituitary surgery, cranial radiation tx, tumor treatment, TBI or subarachnoid hemorrhage,

PA Criteria	Criteria Details
	<p>AND 3. meets one of the following - A. has known perinatal insults, congenital or genetic defects or structural hypothalamic pituitary defects, B. 3 or more pituitary hormone def (ACTH, TSH, LH/FSH, or prolactin, age and gender adjusted IGF1 below the lower limits of the normal reference range AND other causes of low serum IGF-1 have been excluded, C. Neg response to ONE preferred GH stim test (insulin peak response less than or equal to 5 mcg/L, Glucagon peak less than or equal to 3 mcg/L (BMI is less than or equal to 25), less than or equal to 3 and BMI is greater than or equal to 25 and less than or equal to 30 with a high pretest probability of GH deficiency, less than or equal to 1 and BMI is greater than or equal to 25 and less than or equal to 30 with a low pretest probability of GH deficiency or less than or equal to 1 mcg/L (BMI is greater than 30), if insulin and glucagon contraindicated then Arginine test with peak of less than or equal to 0.4 mcg/L, or Macrilen peak less than 2.8 ng/ml AND BMI is less than or equal to 40 AND if a transitional adolescent must be off tx for at least one month before retesting. Cont tx - endocrine must certify not being prescribed for anti-aging or to enhance athletic performance. ISS initial - baseline ht less than the 1.2 percentile or a standard deviation score (SDS) less than -2.25 for age and gender, open epiphyses, does not have CDGP and height velocity is either growth rate (GR) is a. less than 4 cm/yr for pts greater than or equal to 5 or b. growth velocity is less than 10th percentile for age/gender. Cont tx - prescriber confirms response to therapy. CKD initial - CKD defined by abnormal CrCl. Noonan initial - baseline height less than 5th percentile. PW cont tx in adults or adolescents who don't meet child requir - physician certifies not being used for anti-aging or to enhance athletic performance. SHOX initial - SHOX def by chromo analysis, open epiphyses, height less than 3rd percentile for age/gender. SGA initial -baseline ht less than 5th percentile for age/gender and born SGA (birth weight/length that is more than 2 SD below mean for gestational age/gender and didn't have sufficient catch up growth by 2-4 y/o). Cont tx - prescriber confirms response to therapy. Cont Tx for CKD, Noonan, PW in child/adoles, SHOX, and TS - prescriber confirms response to therapy. SBS initial pt receiving specialized nutritional support. Cont tx - 2nd course if pt responded to tx with a decrease in the requirement for specialized nutritional support.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Short bowel syndrome
Part B Prerequisite	No

HERNEXEOS

Products Affected

- HERNEXEOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NON-SMALL CELL LUNG CANCER-all the following (A, B, C, and D): A) Unresectable or metastatic disease, AND B) Human epidermal growth factor receptor 2 (HER2) [ERBB2] activating mutation, AND C) Mutation was detected by an approved test, AND D) Received at least one prior systemic therapy. Note: Examples include checkpoint inhibitors such as Keytruda (pembrolizumab intravenous infusion), Libtayo (cemiplimab-rwlc intravenous infusion), Tecentriq (atezolizumab intravenous infusion), Opdivo (nivolumab intravenous infusion), or Imjudo (tremelimumab-actl intravenous infusion) in combination with chemotherapy (e.g., carboplatin, cisplatin, pemetrexed, paclitaxel, albumin-bound paclitaxel, bevacizumab), chemotherapy alone (e.g., docetaxel, gemcitabine, etoposide, vinorelbine, other chemotherapy noted above).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS - BENZODIAZEPINES

Products Affected

- *clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg*
- *diazepam intensol*
- *diazepam oral solution 5 mg/5 ml (1 mg/ml)*
- *diazepam oral tablet*
- *lorazepam intensol*
- *lorazepam oral tablet 0.5 mg, 1 mg, 2 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Procedure-related sedation = 1mo. All other conditions = 12 months.
Other Criteria	All medically accepted indications other than insomnia, approve if the physician has assessed risk versus benefit in using the High Risk Medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy. Insomnia, may approve lorazepam if the patient has had a trial with two of the following: ramelteon, doxepin 3mg or 6 mg, eszopiclone, zolpidem, or zaleplon and the physician has assessed risk versus benefit in using the HRM in this patient and has confirmed that he/she would still like initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS - BENZTROPINE

Products Affected

- *benztropine oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For all medically-accepted indications, approve if the prescriber confirms he/she has assessed risk versus benefit in prescribing benztropine for the patient and he/she would still like to initiate/continue therapy
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS - CYCLOBENZAPRINE

Products Affected

- *cyclobenzaprine oral tablet 10 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	The physician has assessed risk versus benefit in using this High Risk Medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS - FIRST GENERATION ANTIHISTAMINES

Products Affected

- *hydroxyzine hcl oral tablet*
- *promethazine oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For hydroxyzine hydrochloride, authorize use without a previous drug trial for all FDA-approved indications other than anxiety. Approve hydroxyzine hydrochloride if the patient has tried at least two other FDA-approved products for the management of anxiety. Prior to approval of promethazine and hydroxyzine, approve if the physician must have assessed risk versus benefit in prescribing the requested HRM for the patient and must confirm that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS - PHENOBARBITAL

Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia.
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the treatment of seizures, approve only if the patient is currently taking phenobarbital.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATIONS- ESTROGENS

Products Affected

- *abigale*
- *abigale lo*
- *dotti*
- *estradiol oral*
- *estradiol transdermal patch semiweekly*
- *estradiol transdermal patch weekly*
- *estradiol-norethindrone acet*
- *fyavolv*
- *jinteli*
- *lyllana*
- *mimvey*
- *norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Previous medication use
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	For the treatment of Vulvar Vaginal Atrophy, approve if the patient has had a trial of one of the following for vulvar vaginal atrophy (brand or generic): Estradiol Vaginal Cream, Imvexxy, Premarin Vaginal Cream or estradiol valerate injection. For prophylaxis of Postmenopausal Osteoporosis, approve if the patient has had a trial of one of the following (brand or generic): alendronate, ibandronate, risedronate or Raloxifene. The physician has assessed risk versus benefit in using this High Risk medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

IBRANCE

Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast cancer - approve recurrent or metastatic, hormone receptor positive (HR+) [i.e., estrogen receptor positive- (ER+) and/or progesterone receptor positive (PR+)] disease, and HER2-negative breast cancer when the pt meets ONE of the following 1. Pt is postmenopausal and Ibrance will be used in combination with anastrozole, exemestane, or letrozole 2. pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonists, or has had surgical bilateral oophorectomy, or ovarian irradiation AND meets one of the following conditions: Ibrance will be used in combination with anastrozole, exemestane, or letrozole or Ibrance will be used in combination with fulvestrant 3. pt is a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) who is receiving GnRH analog AND Ibrance will be used in combination with anastrozole, exemestane or letrozole or Ibrance will be used in combination with fulvestrant 4. Pt is postmenopausal and Ibrance will be used in combination with fulvestrant. Liposarcoma-approve if the patient has well-differentiated/dedifferentiated liposarcoma (WD-DDLS).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Liposarcoma

PA Criteria	Criteria Details
Part B Prerequisite	No

IBTROZI

Products Affected

- IBTROZI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NON-SMALL CELL LUNG CANCER-locally advanced or metastatic disease and ROS1-positive non-small cell lung cancer as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ICATIBANT

Products Affected

- *icatibant*
- *sajazir*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by, or in consultation with, an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Hereditary Angioedema (HAE) Due to C1 Inhibitor (C1-INH) Deficiency [Type I or Type II] - Treatment of Acute Attacks, Initial Therapy-the patient has HAE type I or type II as confirmed by the following diagnostic criteria (i and ii): i. the patient has low levels of functional C1-INH protein (less than 50 percent of normal) at baseline, as defined by the laboratory reference values AND ii. the patient has lower than normal serum C4 levels at baseline, as defined by the laboratory reference values. Patients who have treated previous acute HAE attacks with icatibant - the patient has treated previous acute HAE type I or type II attacks with icatibant AND according to the prescribing physician, the patient has had a favorable clinical response (e.g., decrease in the duration of HAE attacks, quick onset of symptom relief, complete resolution of symptoms, decrease in HAE acute attack frequency or severity) with icatibant treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ICLUSIG

Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis the Philadelphia chromosome (Ph) status of the leukemia must be reported. T315I status
Age Restrictions	All indications except ALL - 18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Acute lymphoblastic leukemia, Philadelphia chromosome positive or ABL-class translocation-approve. Chronic myeloid leukemia-approve. GIST - approve if the patient tried all of the following therapies first to align with NCCN recommendations which include: Imatinib or Ayyvakit (avapritinib tablets), AND Sunitinib or Sprycel (dasatinib tablets), AND Stivarga (regorafenib tablets), AND Qinlock (ripretinib tablets). Myeloid/Lymphoid Neoplasms with Eosinophilia - approve if the tumor has ABL1 rearrangement or FGFR1 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Gastrointestinal Stromal Tumor, Myeloid/Lymphoid Neoplasms with Eosinophilia
Part B Prerequisite	No

IDHIFA

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	IDH2-mutation status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	AML - approve if the patient is IDH2-mutation status positive as detected by an approved test
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

IMATINIB

Products Affected

- *imatinib oral tablet 100 mg, 400 mg*
- IMKELDI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported.
Age Restrictions	ASM, DFSP, HES, MDS/MPD/Myeloid/Lymphoid Neoplasms/Kaposi Sarcoma/Cutaneous Melanoma-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For ALL-approve for Ph-positive or ABL-class translocation ALL. CML-approve for Ph-positive or BCR::ABL1-mutation positive CML. Kaposi's Sarcoma-approve if the patient has tried at least one regimen AND has relapsed or refractory disease. Pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT)-patient has tried Turalio or Romvimza or according to the prescriber, the patient cannot take Turalio or Romvimza. Myelodysplastic/myeloproliferative disease-approve if the condition is associated with platelet-derived growth factor receptor (PDGFR) gene rearrangements. Graft versus host disease, chronic-approve if the patient has tried at least one conventional systemic treatment (e.g., imbruvica). Cutaneous melanoma-approve if the patient has an activating KIT mutation, metastatic or unresectable melanoma, and has tried at least one systemic regimen. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an ABL1 rearrangement or an FIP1L1-PDGFR α or PDGFR β rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chordoma, desmoid tumors (aggressive fibromatosis), cKit positive metastatic or unresectable cutaneous melanoma, Kaposi's Sarcoma and pigmented Villonodular Synovitis/Tenosynovial Giant Cell Tumor, myeloid/lymphoid neoplasms with eosinophilia, GVHD, chronic.

PA Criteria	Criteria Details
Part B Prerequisite	No

IMBRUVICA

Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	GVHD-1 year and older, other-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	CLL- Approve. GVHD-Approve if the patient has tried one conventional systemic treatment for graft versus host disease (e.g., corticosteroids [methylprednisolone, prednisone], imatinib, low-dose methotrexate, sirolimus, mycophenolate mofetil, Jakafi [ruxolitinib tablets]). B-cell lymphoma-approve if the patient has tried at least one systemic regimen (e.g., cisplatin, cytarabine, rituximab, oxaliplatin, gemcitabine, ifosfamide, carboplatin, etoposide, or rituximab). Central nervous system Lymphoma (primary)- approve if the patient is not a candidate for or is intolerant to high-dose methotrexate OR has tried at least one therapy (e.g., methotrexate, rituximab, vincristine, procarbazine, cytarabine, thiotepa, carmustine, intrathecal methotrexate, cytarabine, or rituximab). Hairy Cell Leukemia - approve if the patient has tried at least two systemic regimens (cladribine, Nipent [pentostatin injection], rituximab, or Pegasys [peginterferon alfa-2a subcutaneous injection]).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Central Nervous System Lymphoma (Primary), Hairy Cell Leukemia, B-Cell Lymphoma
Part B Prerequisite	No

INBRIJA

Products Affected

- INBRIJA INHALATION CAPSULE,
W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	Asthma, COPD, other chronic underlying lung disease
Required Medical Information	Diagnosis, medications that will be used in combination
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Approve if the patient is currently taking carbidopa-levodopa and is experiencing off episodes.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

INGREZZA

Products Affected

- INGREZZA
- INGREZZA INITIATION PK(TARDIV)
- INGREZZA SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	TD - Prescribed by or in consultation with a neurologist or psychiatrist. Chorea HD - prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Chorea associated with Huntington's Disease- approve if diagnosis is confirmed by genetic testing (for example, an expanded HTT CAG repeat sequence of at least 36).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

INJECTABLE TESTOSTERONE PRODUCTS

Products Affected

- *testosterone cypionate*
- *testosterone enanthate*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, lab results
Age Restrictions	Delayed puberty or induction of puberty in males-14 years and older, 12 years and older (testosterone cypionate)
Prescriber Restrictions	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of transgender patients.
Coverage Duration	Delayed puberty or induction of puberty in males-6 months, all others-12 months
Other Criteria	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. Delayed puberty or induction of puberty in males - Approve testosterone enanthate or testosterone cypionate. Breast cancer in females-approve testosterone enanthate. Male is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression. Female is defined as an individual with the biological traits of a woman, regardless of the individual's gender identity or gender expression. Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-

PA Criteria	Criteria Details
	Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-approve. Note: For a patient who has undergone gender reassignment, use this FTM criterion for hypogonadism indication.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization).
Part B Prerequisite	No

INLYTA

Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Advanced Renal cell carcinoma-approve. Differentiated thyroid cancer, approve if patient is refractory to radioactive iodine therapy. Soft tissue sarcoma-approve if the patient has alveolar soft part sarcoma and the medication will be used in combination with Keytruda (pembrolizumab).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (i.e., papillary, follicular, and oncocytic) Thyroid Carcinoma, Soft tissue sarcoma
Part B Prerequisite	No

INPEFA

Products Affected

- INPEFA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Heart Failure, to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit-approve. Type 2 Diabetes, to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit-approve if the patient has chronic kidney disease AND has one or more cardiovascular risk factor(s). Note: Patients with heart failure should be reviewed under criteria for Heart Failure.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

INQOVI

Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myelodysplastic Syndrome/Myeloproliferative Neoplasm Overlap Neoplasms
Part B Prerequisite	No

INREBIC

Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Myelofibrosis (MF), including Primary MF, Post-Polycythemia Vera MF, and Post-Essential Thrombocythemia MF-approve if the patient has intermediate-2 or high-risk disease. Myeloid/Lymphoid Neoplasms with Eosinophilia-approve if the tumor has a JAK2 rearrangement. Accelerated or blast phase myeloproliferative neoplasm- approve if the patient has at least one disease-related symptom (examples: fatigue, fever, night sweats, weight loss, abdominal discomfort, splenomegaly, thrombocytosis, or leukocytosis).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myeloid/Lymphoid Neoplasms with Eosinophilia, accelerated or blast phase myeloproliferative neoplasm
Part B Prerequisite	No

ITOVEBI

Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	BREAST CANCER (all of A, B, C, D, E and F): A. Patient meets ONE of the following (i or ii): i. Patient is a postmenopausal female, OR ii. Patient meets BOTH of the following (a and b): a. Patient is a pre/perimenopausal female or a male, AND b. Patient is receiving a gonadotropin-releasing hormone (GnRH) agonist OR had surgical bilateral oophorectomy or ovarian irradiation (female) or orchiectomy (male), Note: Examples of a GnRH agonist include leuprolide acetate, leuprolide acetate intramuscular injection, triptorelin pamoate intramuscular injection, goserelin acetate subcutaneous injection. AND B. Patient has locally advanced or metastatic hormone receptor (HR)-positive disease, AND C. Patient has human epidermal growth factor receptor 2 (HER2)-negative disease, AND D. Patient has PIK3CA-mutated breast cancer as detected by an approved test, AND E. Patient meets (i or ii): i) has disease progression while on adjuvant endocrine therapy or ii) had disease recurrence within 12 months after completing adjuvant endocrine therapy, Note: Examples of endocrine therapy include tamoxifen, anastrozole, letrozole, exemestane, toremifene. AND F. The medication will be used in combination with palbociclib capsules/tablets and fulvestrant injection.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

IVERMECTIN (ORAL)

Products Affected

- *ivermectin oral tablet 3 mg, 6 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	30 days
Other Criteria	Pediculosis-approve if the patient has infection caused by pediculus humanus capitis (head lice), pediculus humanus corporis (body lice), or has pediculosis pubis caused by phthirus pubis (pubic lice). Scabies-approve if the patient has classic scabies, treatment resistant scabies, is unable to tolerate topical treatment, has crusted scabies or is using ivermectin tablets for prevention and/or control of scabies.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Ascariasis, Enterobiasis (pinworm infection), Hookworm-related cutaneous larva migrans, Mansonella ozzardi infection, Mansonella streptocerca infection, Pediculosis, Scabies, Trichuriasis, Wuchereria bancrofti infection
Part B Prerequisite	No

IVIG

Products Affected

- PRIVIGEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Part B versus D determination per CMS guidance to establish if drug used for PID in pt's home.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

IWILFIN

Products Affected

- IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Neuroblastoma-Approve if the patient meets the following (A, B and C): A) Patient has high-risk disease, AND B) The medication is being used to reduce the risk of relapse, AND C) Patient has had at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy. Note: Examples of anti-GD2 immunotherapy includes Unituxin (dinutuximab intravenous infusion).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

JAKAFI

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	ALL-1 to 21 years of age, GVHD-12 and older, MF/PV/accelerated or blast phase MPN/CMML-2/essential thrombo/myeloid/lymphoid neoplasm/T-cell Lymphoma-18 and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For polycythemia vera patients must have tried hydroxyurea or peginterferon alfa-2a or Besremi (ropoferon alfa-2b-njft subcutaneous injection). ALL-approve if the mutation/pathway is Janus associated kinase (JAK)-related. GVHD, chronic-approve if the patient has tried one conventional systemic treatment for graft versus host disease (for example: prednisone, ibrutinib capsules/tablets). GVHD, acute-approve if the patient has tried one systemic corticosteroid. Atypical chronic myeloid leukemia-approve if the patient has a CSF3R mutation or a janus associated kinase 2 (JAK2) mutation. Chronic monomyelocytic leukemia-2 (CMML-2)-approve if the patient is also receiving a hypomethylating agent. Essential thrombocythemia-approve if the patient has tried hydroxyurea, peginterferon alfa-2a or anagrelide. Myeloid/lymphoid neoplasms-approve if the patient has eosinophilia and the tumor has a janus associated kinase 2 (JAK2) rearrangement. T-Cell Lymphoma - approve if pt has (A or B): A) peripheral T-cell lymphoma or B) meets (i and ii): i) pt has T-cell prolymphocytic leukemia, T-cell large granular lymphocytic leukemia, hepatosplenic T-cell lymphoma, or breast implant-associated anaplastic large cell lymphoma and ii) pt has tried at least one systemic regimen. Accelerated or blast phase myeloproliferative neoplasm-approve if pt has at least one disease-related symptom (examples: fatigue, fever, night sweats,

PA Criteria	Criteria Details
	weight loss, abdominal discomfort, splenomegaly, thrombocytosis, or leukocytosis).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Acute lymphoblastic leukemia, atypical chronic myeloid leukemia, chronic monomyelocytic leukemia-2 (CMML-2), essential thrombocythemia, myeloid/lymphoid neoplasms, T-Cell lymphoma, accelerated or blast phase myeloproliferative neoplasm
Part B Prerequisite	No

JAYPIRCA

Products Affected

- JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	<p>Mantle cell lymphoma-approve if the patient has tried at least one systemic regimen or patient is not a candidate for a systemic regimen (i.e., an elderly patient who is frail), AND the patient has tried one Bruton tyrosine kinase inhibitor (BTK) for mantle cell lymphoma. Note: Examples of a systemic regimen contain one or more of the following products: rituximab, dexamethasone, cytarabine, carboplatin, cisplatin, oxaliplatin, cyclophosphamide, doxorubicin, vincristine, prednisone, methotrexate, bendamustine, Velcade (bortezomib intravenous or subcutaneous injection), lenalidomide, gemcitabine, and Venclexta (venetoclax tablets). Note: Examples of BTK inhibitors indicated for mantle cell lymphoma include Brukinsa (zanubrutinib capsules), Calquence (acalabrutinib capsules), and Imbruvica (ibrutinib capsules, tablets, and oral suspension).</p> <p>CLL/SLL-patient meets (A or B): A) patient has resistance or intolerance to Imbruvica (ibrutinib tablets, capsules, or oral solution), Calquence (acalabrutinib tablets), or Brukinsa (zanubrutinib capsules) or B) patient has relapsed or refractory disease and has tried a Bruton tyrosine kinase (BTK) inhibitor and Venclexta (venetoclax tablet)-based regimen.</p> <p>Examples of BTK inhibitor include: Imbruvica (ibrutinib tablets, capsules, or oral solution), Calquence (acalabrutinib tablets), or Brukinsa (zanubrutinib capsules). Richter's Transformation to DLBCL- pt has tried at least one chemotherapy regimen or is not a candidate for a chemotherapy</p>

PA Criteria	Criteria Details
	regimen. Marginal Zone Lymphoma - approve if pt has tried at least one Bruton tyrosine kinase inhibitor.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Richter's Transformation to Diffuse Large B-Cell Lymphoma, Marginal Zone Lymphoma
Part B Prerequisite	No

JYNARQUE

Products Affected

- *tolvaptan (polycys kidney dis) oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	Patient is currently receiving another tolvaptan product.
Required Medical Information	Diagnosis, renal function
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist
Coverage Duration	1 year (initial and continuation)
Other Criteria	Approve if the patient has rapidly-progressing autosomal dominant polycystic kidney disease (ADPKD) (e.g., reduced or declining renal function, high or increasing total kidney volume [height adjusted]), according to the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Orkambi, Trikafta or Symdeko
Required Medical Information	N/A
Age Restrictions	1 month of age and older
Prescriber Restrictions	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	1 year
Other Criteria	CF - must meet A, B, and C: A) pt must have one mutation in the CFTR gene that is considered to be pathogenic or likely pathogenic B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KERENDIA

Products Affected

- KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with spironolactone or eplerenone
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (initial and continuation therapy)
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Diabetic kidney disease-initial-approve if the patient meets the following criteria (i, ii and iii)i. Patient has a diagnosis of type 2 diabetes AND ii. Patient meets one of the following (a or b): a) Patient is currently receiving a maximally tolerated labeled dosage of an angiotensin converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) OR b) According to the prescriber, patient has a contraindication or has experienced significant intolerance to ACE inhibitor and ARB therapy AND iii. At baseline (prior to the initiation of Kerendia), patient meets all of the following (a, b, and c): a) Estimated glomerular filtration rate greater than or equal to 25 mL/min/1.73 m ² AND b) Urine albumin-to-creatinine ratio greater than or equal to 30 mg/g AND c) Serum potassium level less than or equal to 5.0 mEq/L. Diabetic kidney disease-continuation-approve if the patient meets the following criteria (i and ii): i. Patient has a diagnosis of type 2 diabetes AND ii. Patient meets one of the following (a or b): a) Patient is currently receiving a maximally tolerated angiotensin converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) OR b) According to the prescriber, patient has a contraindication or has experienced significant intolerance to ACE inhibitor and ARB therapy. Heart failure-initial-Pt meets all (i, ii, and iii): i) has left ventricular ejection fraction of at least 40 percent AND ii) tried one of the following SGLT-2 inhibitors: Farxiga (dapagliflozin tabs, authorized generic) Inpefa (sotagliflozin tabs), or Jardiance (empagliflozin tabs) OR has contraindication or has experienced

PA Criteria	Criteria Details
	significant intolerance to SGLT-2 inhibitors AND iii) at baseline (prior to initiation of Kerendia), meets all (a and b): a) estimated glomerular filtration rate of at least 25 mL/min/1.73m ² AND b) serum potassium level of less than or equal to 5.0 mEq/L. Heart failure-continuation-Pt meets all (i and ii): i) has left ventricular ejection fraction of at least 40 percent AND ii) tried one of the following SGLT-2 inhibitors: Farxiga (dapagliflozin tabs, authorized generic) Inpefa or (sotagliflozin tabs), or Jardiance (empagliflozin tabs) or has contraindication or has experienced significant intolerance to SGTL-2 inhibitors.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KESIMPTA

Products Affected

- KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS)
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include, clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease.
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KISQALI

Products Affected

- KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast cancer - approve for hormone receptor positive (HR+) [i.e., estrogen receptor positive (ER+) and/or progesterone receptor positive (PR+)] disease, and HER2-negative early (stage II or III), recurrent, or metastatic breast cancer [for early breast cancer must be adjuvant treatment and high risk of recurrence] when the pt meets ONE of the following 1. Pt is postmenopausal and Kisqali will be used in combination with anastrozole, exemestane, or letrozole 2. pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonist, or has had surgical bilateral oophorectomy, or ovarian irradiation AND Kisqali will be used in combination with anastrozole, exemestane, or letrozole 3. pt is a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) who is receiving GnRH analog AND Kisqali will be used in combination with anastrozole, exemestane or letrozole. 4. Patient is postmenopausal, pre/perimenopausal (patient receiving ovarian suppression/ablation with a GnRH agonist or has had surgical bilateral oophorectomy or ovarian irradiation) or a man, and Kisqali (not Co-Pack) will be used in combination with fulvestrant. If the request is for Kisqali Femara, patients do not need to use in combination with anastrozole, exemestane or letrozole. Endometrial cancer - approve if pt meets all of (A,

PA Criteria	Criteria Details
	B and C): A) pt has recurrent or metastatic disease, and B) has estrogen receptor (ER)-positive tumors, and C) if request is for Kisqali (not Co-Pack), Kisqali will be used in combination with letrozole.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Endometrial cancer
Part B Prerequisite	No

KORLYM

Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior surgeries
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of Cushing's syndrome
Coverage Duration	1 year
Other Criteria	Endogenous Cushing's Syndrome-Approve if mifepristone is being used to control hyperglycemia secondary to hypercortisolism in patients who have type 2 diabetes mellitus or glucose intolerance AND pt meets (i, ii or iii): i) patient is not a candidate for surgery or surgery has not been curative, or (ii) patient is awaiting surgery for endogenous Cushing's Syndrome or (iii) patient is awaiting therapeutic response after radiotherapy for endogenous Cushing's Syndrome.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KOSELUGO

Products Affected

- KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	<p>Neurofibromatosis Type 1-approve if prior to starting Koselugo, the patient has symptomatic, inoperable plexiform neurofibromas and if the patient is 2 to 18 years old OR if the patient is 19 years or older if the patient started on therapy with Koselugo prior to becoming 19. Circumscribed Glioma-approve if the patient has recurrent, refractory or progressive disease AND the tumor is BRAF fusion positive OR BRAF V600E activating mutation positive OR patient has neurofibromatosis type 1 mutated glioma AND this medication will be used as a single agent AND the patient is 3-21 years of age OR is greater 21 and has been previously started on therapy with Koselugo prior to becoming 21 years of age. Langerhans Cell Histiocytosis- approve if the patient meets the following criteria (A and B): A) Patient meets one of the following (i, ii, iii, or iv): i. Patient meets both of the following (a and b): a) Patient has multisystem Langerhans cell histiocytosis, AND b) Patient has symptomatic disease or impending organ dysfunction, OR ii. Patient has single system lung Langerhans cell histiocytosis, OR iii. Patient meets all of the following (a, b, and c): a) Patient has single system bone disease, AND b) Patient has not responded to treatment with a bisphosphonate, AND c) Patient has more than 2 bone lesions, OR iv. Patient has central nervous system disease, AND B) The medication is used as a single agent.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Circumscribed Glioma, Langerhans Cell Histiocytosis
Part B Prerequisite	No

KRAZATI

Products Affected

- KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-approve if (A and B): A) the patient has KRAS G12C-mutated locally advanced or metastatic NSCLC, as determined by an approved test AND B) patient meets either (i or ii): i) has been previously treated with at least one systemic regimen [Examples of systemic regimens include those containing one or more of the following products: Keytruda (pembrolizumab intravenous infusion), Opdivo (nivolumab intravenous infusion), Tecentriq (atezolizumab intravenous infusion), Alimta (pemetrexed intravenous infusion), Yervoy (ipilimumab intravenous infusion), Abraxane (albumin-bound paclitaxel intravenous infusion), bevacizumab, cisplatin, carboplatin, docetaxel, gemcitabine, paclitaxel, vinorelbine.] or ii) patient has brain metastases. Colon or Rectal Cancer- approve if pt has unresectable, advanced, or metastatic disease AND pt has KRAS G12C mutation-positive disease AND medication is prescribed as part of a combination regimen or the patient is unable to tolerate combination therapy AND pt has previously received a chemotherapy regimen for colon or rectal cancer. Ampullary adenocarcinoma-approve if (A, B and C): A) metastatic disease, B) KRAS G12C mutation-positive disease, and C) will be used as subsequent therapy. Biliary tract cancer- approve if (A, B and C): A) unresectable or metastatic disease, B) KRAS G12C mutation-positive disease, and C) previously treated with at least one systemic regimen. Pancreatic adenocarcinoma- approve if (A and B): A) KRAS G12C mutation-positive

PA Criteria	Criteria Details
	disease, and B) either (i or ii): (i) locally advanced or metastatic disease and previously treated with at least one systemic regimen, or (ii) recurrent disease after resection. Small bowel adenocarcinoma- approve if (A, B and C): A) advanced or metastatic disease, B) KRAS G12C mutation-positive disease, and C) will be used as subsequent therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Ampullary adenocarcinoma, biliary tract cancer, pancreatic adenocarcinoma, small bowel adenocarcinoma
Part B Prerequisite	No

LAPATINIB

Products Affected

- *lapatinib*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which lapatinib is being used. Metastatic breast cancer, HER2 status or hormone receptor (HR) status.
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	HER2-positive recurrent or metastatic breast cancer, approve if lapatinib will be used in combination with capecitabine OR trastuzumab and the patient has tried at least two prior anti-HER2 based regimens OR the medication will be used in combination with an aromatase inhibitor and the patient has HR+ disease and the patient is a postmenopausal woman or the patient is premenopausal or perimenopausal woman and is receiving ovarian suppression/ablation with a GnRH agonist, surgical bilateral oophorectomy or ovarian irradiation OR the patient is a man and is receiving a GnRH analog. Colon or rectal cancer-approve if the patient has unresectable advanced or metastatic disease that is human epidermal receptor 2 (HER2) amplified and with wild-type RAS and BRAF disease and the patient has tried at least one chemotherapy regimen or is not a candidate for intensive therapy and the medication is used in combination with trastuzumab (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan) and the patient has not been previously treated with a HER2-inhibitor. Bone Cancer-approve if the patient has recurrent chordoma and if the patient has epidermal growth-factor receptor (EGFR)-positive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Bone cancer-chordoma, colon or rectal cancer, breast cancer in pre/peri-menopausal women and men

PA Criteria	Criteria Details
Part B Prerequisite	Yes

LAZCLUZE

Products Affected

- LAZCLUZE ORAL TABLET 240 MG,
80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NON-SMALL CELL LUNG CANCER-ALL of the following (A, B, C, and D): A. Locally advanced or metastatic disease, AND B. Epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations, as detected by an approved test, AND C. Used in combination with Rybrevant, AND D. Used as first-line treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LEDIPASVIR/SOFOSBUVIR

Products Affected

- LEDIPASVIR-SOFOSBUVIR

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with other direct acting antivirals, excluding ribavirin
Required Medical Information	Diagnosis
Age Restrictions	3 years or older
Prescriber Restrictions	Prescribed by or in consultation w/ GI, hepatologist, ID, or a liver transplant MD
Coverage Duration	Criteria will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	Criteria will be applied consistent with current AASLD/IDSA guidance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance
Part B Prerequisite	No

LENALIDOMIDE

Products Affected

- *lenalidomide*
- REVLIMID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and previous therapies or drug regimens tried.
Age Restrictions	18 years and older (except Kaposi's Sarcoma, Castleman's Disease, CNS Lymphoma)
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Follicular lymphoma-approve if the patient is using lenalidomide in combination with rituximab or has tried at least one prior therapy. MCL-approve -if the patient is using lenalidomide in combination with rituximab or has tried at least two other therapies or therapeutic regimens. MZL-approve if the patient is using lenalidomide in combination with rituximab or has tried at least one other therapy or therapeutic regimen. Multiple myeloma-approve. MDS-approve if the patient meets one of the following: 1) Pt has symptomatic anemia, OR 2) Pt has transfusion-dependent anemia, OR 3) Pt has anemia that is not controlled with an erythroid stimulating agent (eg, EpoGen, Procrit [epoetin alfa injection], Aranesp [darbepoetin alfa injection]). B-cell-lymphoma (other)-approve if the pt has tried at least one prior therapy. Myelofibrosis-approve if according to the prescriber the patient has anemia. Peripheral T-Cell Lymphoma or T-Cell Leukemia/Lymphoma-approve. CNS lymphoma-approve if according to the prescriber the patient has relapsed or refractory disease. Hodgkin lymphoma, classical-approve if the patient has tried at least three other regimens. Castleman's disease-approve if the patient has relapsed/refractory or progressive disease. Kaposi's Sarcoma-approve if the patient has tried at least one regimen or therapy and the patient has relapsed or refractory disease. Systemic light chain amyloidosis-approve if lenalidomide is used in combination with dexamethasone. Histiocytic

PA Criteria	Criteria Details
	neoplasms-approve if the patient has Langerhans cell histiocytosis with single-system multifocal skin disease or Rosai-Dorfman disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Systemic Amyloidosis Light Chain, Diffuse Large B Cell Lymphoma (Non-Hodgkin's Lymphoma), Myelofibrosis. Castleman's Disease, Hodgkin lymphoma (Classical), Peripheral T-Cell Lymphoma, T-Cell Leukemia/Lymphoma, Central nervous system Lymphoma, Kaposi's sarcoma, histiocytic neoplasms.
Part B Prerequisite	No

LENVIMA

Products Affected

- LENVIMA ORAL CAPSULE 10 MG/DAY (10 MG X 1), 12 MG/DAY (4 MG X 3), 14 MG/DAY(10 MG X 1-4 MG X 1), 18 MG/DAY (10 MG X 1-4 MG X2), 20 MG/DAY (10 MG X 2), 24 MG/DAY(10 MG X 2-4 MG X 1), 4 MG, 8 MG/DAY (4 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	DTC - must be refractory to radioactive iodine treatment for approval. RCC - approve if the pt meets ALL of the following criteria: 1) pt has advanced disease AND if the patient meets i or ii-i. Lenvima is being used in combination with Keytruda OR ii. Lenvima is used in combination with Afinitor/Afinitor Disperz and the patient meets a or b-a. Patient has clear cell histology and patient has tried one antiangiogenic therapy or b. patient has non-clear cell histology. MTC-approve if the patient has tried at least one systemic therapy. Endometrial Carcinoma-Approve if the patient meets the following criteria (A, B, C, and D): A) The patient has advanced endometrial carcinoma that is mismatch repair proficient (pMMR) or not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) AND B) The medication is used in combination with Keytruda (pembrolizumab for intravenous injection) AND C) the disease has progressed on at least one prior systemic therapy AND D) The patient is not a candidate for curative surgery or radiation. HCC-approve if the patient has unresectable or metastatic disease. Thymic carcinoma-approve if the patient has tried at least one chemotherapy regimen. Melanoma - approve if the patient has unresectable or metastatic melanoma AND the medication will be used in combination with Keytruda (pembrolizumab intravenous injection) AND the patient has disease progression on anti-

PA Criteria	Criteria Details
	programmed death receptor-1 (PD-1)/programmed death-ligand 1 (PD-L1)-based therapy. Anaplastic thyroid carcinoma-approve if the medication is used in combination with Keytruda (pembrolizumab intravenous infusion).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Medullary Thyroid Carcinoma (MTC), thymic carcinoma, Melanoma, Anaplastic thyroid carcinoma
Part B Prerequisite	No

LIDOCAINE PATCH

Products Affected

- *lidocaine topical adhesive patch, medicated 5 %*
- *lidocan iii*
- *tridacaine ii*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Diabetic neuropathic pain, chronic back pain
Part B Prerequisite	No

LIVTENCITY

Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with ganciclovir or valganciclovir
Required Medical Information	Diagnosis
Age Restrictions	12 years and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, infectious diseases specialist, oncologist, or a physician affiliated with a transplant center.
Coverage Duration	2 months
Other Criteria	Cytomegalovirus Infection, Treatment-approve if the patient meets the following criteria (A, B, and C): A) Patient weighs greater than or equal to 35 kg, AND B) Patient is post-transplant, AND Note: This includes patients who are post hematopoietic stem cell transplant or solid organ transplant. C) Patient has cytomegalovirus infection/disease that is refractory to treatment with at least one of the following: cidofovir, foscarnet, ganciclovir, or valganciclovir or patient has a significant intolerance to ganciclovir or valganciclovir.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LONG ACTING OPIOIDS

Products Affected

- *BELBUCA*
- *buprenorphine transdermal patch*
- *hydromorphone oral tablet extended release 24 hr*
- *methadone oral solution 10 mg/5 ml, 5 mg/5 ml*
- *methadone oral tablet 10 mg, 5 mg*
- *morphine oral tablet extended release*
- *OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG*

PA Criteria	Criteria Details
Exclusion Criteria	Acute (ie, non-chronic) pain
Required Medical Information	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been tried and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, in hospice, sickle cell disease or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

LONSURF

Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Gastric or (Gas)troesophageal Junction Adenocarcinoma, approve if the patient has been previously treated with at least two chemotherapy regimens for gastric or gastroesophageal junction adenocarcinoma. Colon and rectal cancer-approve per labeling if the patient has been previously treated with a fluoropyrimidine, oxaliplatin, irinotecan and if the patient's tumor or metastases are wild-type RAS (KRAS wild type and NRAS wild type) they must also try Erbitux or Vectibix.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LORBRENA

Products Affected

- LORBRENA ORAL TABLET 100 MG,
25 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, ALK status, ROS1 status, previous therapies
Age Restrictions	Pediatric Diffuse High-Grade Glioma- less than 18 years old, All others- 18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Erdheim Chester disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. Inflammatory myofibroblastic tumor (IMT)-approve if the patient has IMT with ALK translocation. NSCLC - Approve if the patient has ALK-positive advanced or metastatic NSCLC, as detected by an approved test. NSCLC-ROS1 Rearrangement-Positive, advanced or metastatic NSCLC-approve if the patient has tried at least one of crizotinib, entrectinib or ceritinib. Large B-Cell Lymphoma- approve if ALK-positive disease and disease is relapsed or refractory. Pediatric Diffuse High-Grade Glioma- approve if ALK-positive disease and (i or ii): i) used as adjuvant therapy, or ii) used for recurrent or progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-small cell lung cancer (NSCLC)-ROS1 Rearrangement-Positive, Erdheim Chester Disease, Inflammatory Myofibroblastic Tumor (IMT), Large B-Cell Lymphoma, Pediatric Diffuse High-Grade Glioma
Part B Prerequisite	No

LUMAKRAS

Products Affected

- LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-approve if the patient has KRAS G12C-mutated locally advanced or metastatic NSCLC, as determined by an FDA-approved test AND has been previously treated with at least one systemic regimen. Ampullary adenocarcinoma - approve if pt has KRAS G12C-mutated disease as determined by an approved test AND this is used as subsequent therapy. Colon or rectal cancer - approve if pt meets all (A, B, C and D): A) unresectable, advanced, or metastatic disease, and B) KRAS G12C mutation-positive disease, and C) medication is prescribed as part of a combination regimen for colon or rectal cancer [Ex: Lumakras plus cetuximab or panitumumab] or patient is unable to tolerate combination therapy, and D) previously received a chemotherapy regimen for colon or rectal cancer. Pancreatic Adenocarcinoma- approve if patient has KRAS G12C-mutated disease, as determined by an approved test AND either (i or ii): (i) patient has locally advanced or metastatic disease and has been previously treated with at least one systemic regimen OR (ii) patient has recurrent disease after resection.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pancreatic Adenocarcinoma, Ampullary Adenocarcinoma

PA Criteria	Criteria Details
Part B Prerequisite	No

LUPRON DEPOT

Products Affected

- LUPRON DEPOT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Premenstrual disorders - 18 years and older
Prescriber Restrictions	Prostate cancer-prescribed by/consultation with oncologist or urologist. Other cancer diagnosis- prescribed by/consultation with an oncologist. Gender dysphoria/reassignment- prescribed by/consultation with endocrinologist or physician who specializes in treatment of transgender patients
Coverage Duration	uterine leiomyomata - 3 months, abnormal uterine bleeding - 6 months, all others - 12 months
Other Criteria	Endometriosis-approve if the pt has tried one of the following, unless contraindicated: a contraceptive, an oral progesterone or depo-medroxyprogesterone injection. An exception can be made if the pt has previously tried a gonadotropin-releasing hormone [GnRH] agonist (e.g. Lupron Depot) or antagonist (e.g. Orilissa). Head and neck cancer-salivary gland tumor- approve if pt has recurrent, unresectable, or metastatic disease AND androgen receptor-positive disease. Premenstrual disorders including PMS and PMDD- approve if pt has severe refractory premenstrual symptoms AND pt has tried an SSRI or combined oral contraceptive. Prostate cancer - for patients new to therapy requesting Lupron Depot 7.5mg, patients are required to try Orgovyx or Eligard prior to approval.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	abnormal uterine bleeding, breast cancer, gender dysphoria/gender reassignment, head and neck cancer-salivary gland tumors, ovarian cancer including fallopian tube and primary peritoneal cancers, premenstrual disorders including premenstrual syndrome and premenstrual dysphoric disorder, prophylaxis or treatment of uterine bleeding or menstrual

PA Criteria	Criteria Details
	suppression in pts with hematologic malignancy or undergoing cancer treatment or prior to bone marrow or stem cell transplant, uterine cancer
Part B Prerequisite	No

LYNPARZA

Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Ovarian, Fallopian Tube, or Primary Peritoneal Cancer - Maintenance monotherapy-Approve if the patient has a germline or somatic BRCA mutation-positive disease as confirmed by an approved test AND The patient is in complete or partial response to at least one platinum-based chemotherapy regimen (e.g., carboplatin with gemcitabine, carboplatin with paclitaxel, cisplatin with gemcitabine). Ovarian, fallopian tube, or primary peritoneal cancer-maintenance, combination therapy-approve if the medication is used in combination with bevacizumab, the patient has homologous recombination deficiency (HRD)-positive disease, as confirmed by an approved test and the patient is in complete or partial response to first-line platinum-based chemotherapy regimen. Breast cancer, adjuvant-approve if the patient has germline BRCA mutation-positive, HER2-negative breast cancer and the patient has tried neoadjuvant or adjuvant therapy. Breast cancer, recurrent or metastatic disease-approve if the patient has recurrent or metastatic disease and has (i or ii): i) germline BRCA mutation-positive breast cancer or ii) germline PALB2 mutation-positive breast cancer. Pancreatic Cancer-maintenance therapy-approve if the patient has a germline BRCA mutation-positive metastatic disease and the disease has not progressed on at least 16 weeks of treatment with a first-line platinum-based chemotherapy regimen. Prostate cancer-castration resistant-approve if the patient has metastatic disease, the medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog

PA Criteria	Criteria Details
	or the patient has had a bilateral orchiectomy, and the patient meets either (i or ii): i) the patient has germline or somatic homologous recombination repair (HRR) gene-mutated disease, as confirmed by an approved test and the patient has been previously treated with at least one androgen receptor directed therapy or ii) the patient has a BRCA mutation and the medication is used in combination with abiraterone plus one of prednisone or prednisolone. Uterine Leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Uterine Leiomyosarcoma
Part B Prerequisite	No

LYTGOBI

Products Affected

- LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Cholangiocarcinoma-approve if the patient has unresectable locally advanced or metastatic disease, tumor has fibroblast growth factor receptor 2 (FGFR2) gene fusions or other rearrangements as detected by an approved test and if the patient has been previously treated with at least one systemic regimen. Note: Examples of systemic regimens include gemcitabine + cisplatin, 5-fluorouracil + oxaliplatin or cisplatin, capecitabine + cisplatin or oxaliplatin, gemcitabine + Abraxane (albumin-bound paclitaxel) or capecitabine or oxaliplatin, and gemcitabine + cisplatin + Abraxane.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MAVYRET

Products Affected

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	3 years or older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
Coverage Duration	Chronic HCV- Criteria will be applied consistent with AASLD/IDSA guidance, Acute HCV-8 weeks
Other Criteria	For Chronic Hepatitis C Virus, criteria will be applied consistent with current AASLD/IDSA guidance. For Acute Hepatitis C Virus (HCV) Genotype 1, 2, 3, 4, 5, or 6: Approve if the patient meets (A, B and C): A) Does not have cirrhosis OR has compensated cirrhosis AND B) has quantifiable HCV RNA AND C) ONE of the following (i, ii, or iii): i. conversion of negative to positive results in anti-HCV antibody, HCV RNA, and/or HCV core antigen OR ii. signs and symptoms of acute hepatitis C virus OR iii. has engaged in a risk behavior for HCV infection within the prior 6 months.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance
Part B Prerequisite	No

MEGESTROL

Products Affected

- *megestrol oral suspension 400 mg/10 ml*
- *megestrol oral tablet (40 mg/ml), 625 mg/5 ml (125 mg/ml)*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for weight gain for cosmetic reasons.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MEKINIST

Products Affected

- MEKINIST ORAL RECON SOLN
- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Mekinist is being used. For melanoma, thyroid cancer and NSCLC must have documentation of BRAF V600 mutations
Age Restrictions	1 year and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Melanoma must be used in patients with BRAF V600 mutation, and patient has unresectable, advanced (including Stage III or Stage IV disease), or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC requires BRAF V600E Mutation and use in combination with Tafinlar. Thyroid cancer, anaplastic-patient has locally advanced or metastatic anaplastic disease AND Mekinist will be taken in combination with Tafinlar, unless intolerant AND the patient has BRAF V600-positive disease. Ovarian/fallopian tube/primary peritoneal cancer-approve if the patient has recurrent disease and the medication is used for low-grade serous carcinoma or the patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Tafinlar. Biliary Tract Cancer-approve if the patient has tried at least one systemic chemotherapy regimen, patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Tafinlar. Central Nervous System Cancer-approve if the medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma OR ii. Recurrent or progressive disease for one of the following conditions (a, b, c or d): a) glioma OR b) Isocitrate dehydrogenase-2 (IDH2)-mutant astrocytoma OR

PA Criteria	Criteria Details
	<p>c) Glioblastoma or d) Oligodendrolioma OR iii. Melanoma with brain metastases AND patient has BRAF V600 mutation-positive disease AND medication will be taken in combination with Tafinlar (dabrafenib). Histiocytic neoplasm-approve if patient has Langerhans cell histiocytosis and one of the following (a, b, or c): a) Multisystem disease OR b) Pulmonary disease OR c) Central nervous system lesions OR patient has Erdheim Chester disease or Rosai-Dorfman disease. Metastatic or Solid Tumors-Approve if the patient meets the following (A, B, and C): A) Patient has BRAF V600 mutation-positive disease, AND B) The medication will be taken in combination with Tafinlar (dabrafenib capsules), AND C) Patient has no satisfactory alternative treatment options. Hairy Cell Leukemia, approve if pt has not previously been treated with a BRAF inhibitor therapy and this will be used for relapsed/refractory disease and will be taken in combination with Tafinlar. Small bowel adenocarcinoma, approve if pt has BRAF V600E mutation-positive advanced or metastatic disease and this will be used with Tafinlar AND (i or ii): i) this will be used as initial therapy and pt has received previous FOLFOX/CAPEOX therapy in the adjuvant setting within the past 12 months or has a contraindication, or (ii) this will be used as second-line and subsequent therapy.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Histiocytic Neoplasm, Hairy Cell Leukemia
Part B Prerequisite	No

MEKTOVI

Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, BRAF V600 status, concomitant medications
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation AND Mektovi will be used in combination with Braftovi. Histiocytic neoplasm-approve if the patient has Langerhans cell histiocytosis and one of the following (i, ii, or iii): i. multisystem disease OR, ii. pulmonary disease or, iii. central nervous system lesions. NSCLC-approve if pt has BRAF V600E mutation-positive metastatic disease AND this medication will be taken in combination with Braftovi (encorafenib capsules).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Histiocytic Neoplasms
Part B Prerequisite	No

MEMANTINE

Products Affected

- *memantine oral capsule, sprinkle, er 24hr*
- *memantine oral solution*
- *memantine oral tablet*
- *memantine-donepezil*
- NAMZARIC ORAL CAPSULE,SPRINKLE,ER 24HR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Indication for which memantine is being prescribed.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with mild to moderate vascular dementia.
Part B Prerequisite	No

MODAFINIL/ARMODAFINIL

Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	Excessive daytime sleepiness associated with narcolepsy-prescribed by or in consultation with a sleep specialist physician or neurologist
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Excessive sleepiness associated with Shift Work Sleep Disorder (SWSD) - approve if patient is working at least 5 overnight shifts per month. Adjunctive/augmentation treatment for depression in adults if the patient is concurrently receiving other medication therapy for depression. Excessive daytime sleepiness associated with obstructive sleep apnea/hypoapnea syndrome-approve. Excessive daytime sleepiness associated with Narcolepsy-approve if narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Excessive daytime sleepiness (EDS) associated with myotonic dystrophy (modafinil only). Adjunctive/augmentation for treatment of depression in adults (modafinil only).
Part B Prerequisite	No

MODEYSO

Products Affected

- MODEYSO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	HIGH-GRADE GLIOMA (Note: Examples of high-grade glioma include World Health Organization (WHO) Grade 3 or 4 gliomas, such as diffuse midline glioma or glioblastoma)-all the following (A, B and C): A) Histone 3 (H3) K27M mutation, AND B) Recurrent or progressive disease, AND C) Received at one least prior therapy. Note: Examples of prior therapy include radiation, temozolomide, procarbazine, lomustine, or vincristine.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MYFEMBREE

Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, test results
Age Restrictions	18 years and older
Prescriber Restrictions	Fibroids-Prescribed by or in consultation with an obstetrician-gynecologist or a health care practitioner who specializes in the treatment of women's health
Coverage Duration	24 months of total therapy between Myfembree or Oriahnn
Other Criteria	Uterine Fibroids (Leiomyomas)-approve if the patient is premenopausal (before menopause) and is experiencing heavy menstrual bleeding associated with the uterine fibroids, the uterine fibroids have been confirmed by a pelvic ultrasound, including transvaginal ultrasonography or sonohysterography, hysteroscopy, or magnetic resonance imaging. Endometriosis-approve if the patient is premenopausal and patient has previously tried one of the following (i or ii): i. A contraceptive (e.g., combination oral contraceptives, levonorgestrel-releasing intrauterine systems, a depo-medroxyprogesterone injection), unless contraindicated OR ii. An oral progesterone (e.g., norethindrone tablets), unless contraindicated. Note: An exception to this requirement can be made if the patient has previously used a gonadotropin-releasing hormone agonist (e.g., Lupron Depot [leuprolide depot suspension]) or Orilissa (elagolix tablets).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NAYZILAM

Products Affected

- NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications used at the same time
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiepileptic medication(s).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NERLYNX

Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Stage of cancer, HER2 status, previous or current medications tried
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Adjuvant tx-Approve for 1 year (total), advanced or metastatic disease-1 year
Other Criteria	Breast cancer, adjuvant therapy - approve if the patient meets all of the following criteria: patient will not be using this medication in combination with HER2 antagonists, patient has HER2-positive breast cancer AND Patient has completed one year of adjuvant therapy with trastuzumab OR could not tolerate one year of therapy. Breast cancer, recurrent or metastatic disease-approve if the patient has HER-2 positive breast cancer, Nerlynx will be used in combination with capecitabine and the patient has tried at least two prior anti-HER2 based regimens.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NEXLETOL

Products Affected

- NEXLETOL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history (as specified in the Other Criteria field)
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	<p>HYPERLIPIDEMIA WITH HeFH (both A and B): A) meets (a, b, c, or d): a) untreated LDL-C greater than or equal to 190 mg/dL, b) phenotypic confirmation (Note 1) of HeFH, c) Dutch Lipid Network criteria score greater than 5, or d) Simon Broome criteria met threshold for definite or possible/probable, AND B) meets (a or b): a) tried one high-intensity statin (throughout, see Definition 1 below) and ezetimibe and LDL-C remains 70 mg/dL or higher or b) statin intolerant (throughout, see Definition 2 below).</p> <p>ESTABLISHED CVD (both A and B): A) patient has/had one of the following conditions: prior MI, ACS, angina, CVA or TIA, CAD, PAD, coronary or other arterial revascularization procedure, B) meets (a or b): a) tried one high-intensity statin and ezetimibe and LDL-C remains 55 mg/dL or higher or b) statin intolerant.</p> <p>PRIMARY HYPERLIPIDEMIA (not associated with established CVD or HeFH) [A or B]: A) tried one high-intensity statin and ezetimibe for 8 weeks or longer and LDL-C remains 70 mg/dL or higher or B) statin intolerant.</p> <p>Note 1: Examples include mutations at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK) or low-density lipoprotein receptor adaptor protein (LDLRAP1) gene.</p> <p>Definition 1: High intensity statin defined as atorvastatin greater than or equal to 40 mg daily or rosuvastatin greater than or equal to 20 mg daily.</p> <p>Definition 2: Statin intolerance defined as experiencing statin related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of</p>

PA Criteria	Criteria Details
	atorvastatin and rosuvastatin and during both trials the skeletal-related symptoms resolved upon discontinuation of the statin.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NEXLIZET

Products Affected

- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history (as specified in the Other Criteria field)
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	<p>HYPERLIPIDEMIA WITH HeFH (both A and B): A) meets (a, b, c, or d): a) untreated LDL-C greater than or equal to 190 mg/dL, b) phenotypic confirmation (Note 1) of HeFH, c) Dutch Lipid Network criteria score greater than 5, or d) Simon Broome criteria met threshold for definite or possible/probable, AND B) meets (a or b): a) tried one high-intensity statin (throughout, see Definition 1 below) and LDL-C remains 70 mg/dL or higher or b) statin intolerant (throughout, see Definition 2 below).</p> <p>ESTABLISHED CVD (both A and B): A) patient has/had one of the following conditions: prior MI, ACS, angina, CVA or TIA, CAD, PAD, coronary or other arterial revascularization procedure, B) meets (a or b): a) tried one high-intensity statin and LDL-C remains 55 mg/dL or higher or b) statin intolerant.</p> <p>PRIMARY HYPERLIPIDEMIA (not associated with established CVD or HeFH) [A or B]: A) tried one high-intensity statin for 8 weeks or longer and LDL-C remains 70 mg/dL or higher or B) statin intolerant.</p> <p>Note 1: Examples include mutations at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK) or low-density lipoprotein receptor adaptor protein (LDLRAP1) gene.</p> <p>Definition 1: High intensity statin defined as atorvastatin greater than or equal to 40 mg daily or rosuvastatin greater than or equal to 20 mg daily.</p> <p>Definition 2: Statin intolerance defined as experiencing statin related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of</p>

PA Criteria	Criteria Details
	atorvastatin and rosuvastatin and during both trials the skeletal-related symptoms resolved upon discontinuation of the statin.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NILUTAMIDE

Products Affected

- *nilutamide*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate cancer-approve if nilutamide is used concurrently with a luteinizing hormone-releasing hormone (LHRH) agonist.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NINLARO

Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	MM - be used in combination with dexamethasone and lenalidomide or cyclophosphamide OR pt had received at least ONE previous therapy for multiple myeloma OR the agent will be used following autologous stem cell transplantation (ASCT). Systemic light chain amyloidosis-approve if the patient has tried at least one other regimen for this condition. Waldenstrom Macroglobulinemia/lymphoplasmacytic lymphoma-approve if used in combination with a rituximab product and dexamethasone (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with systemic light chain amyloidosis, Waldenstrom Macroglobulinemia/lymphoplasmacytic lymphoma, Multiple myeloma after previous treatment (either monotherapy or in combination other than lenalidomide/dexamethasone) or stem cell transplant
Part B Prerequisite	Yes

NITISINONE

Products Affected

- *nitisinone*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant therapy with nitisinone products
Required Medical Information	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
Coverage Duration	1 year
Other Criteria	Hereditary Tyrosinemia, Type 1-approve if the prescriber confirms the diagnosis was confirmed by genetic testing confirming biallelic pathogenic/likely pathogenic variants in the FAH gene OR elevated levels of succinylacetone in the serum or urine.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NIVESTYM

Products Affected

- NIVESTYM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Cancer/AML, MDS, ALL-oncologist or a hematologist. Cancer patients receiving BMT and PBPC-prescribed by or in consultation with an oncologist, hematologist, or a physician who specializes in transplantation. Radiation-expertise in acute radiation. SCN-hematologist. HIV/AIDS neutropenia-infectious disease (ID) physician (MD), hematologist, or MD specializing in HIV/AIDS.
Coverage Duration	chemo/SCN/AML-6 mo.HIV/AIDS-4 mo.MDS-3 mo.PBPC,Drug induce A/N,ALL,BMT, Radiation-1 mo
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient meets one of the following conditions: 1)patient is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), 2)patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status, or HIV infection), 3)patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor (eg, Leukine, filgrastim products, pegfilgrastim products) and a reduced dose or frequency of chemotherapy may compromise treatment, or 4)patient has received chemotherapy has febrile neutropenia and has at least one risk factor (eg, sepsis syndrome, aged greater than 65 years, severe neutropenia

PA Criteria	Criteria Details
	[absolute neutrophil count less than 100 cells/mm ³], neutropenia expected to be greater than 10 days in duration, invasive fungal infection).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Neutropenia associated with human immunodeficiency virus (HIV) or acquired immunodeficiency syndrome (AIDS). Treatment of myelodysplastic syndromes (MDS). Drug induced agranulocytosis or neutropenia. Acute lymphocytic leukemia (ALL), Radiation Syndrome (Hematopoietic Syndrome of Acute Radiation Syndrome)
Part B Prerequisite	No

NON-INJECTABLE TESTOSTERONE PRODUCTS

Products Affected

- *testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)*
- *testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram), 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)*
- *testosterone transdermal solution in metered pump w/app*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of primary hypogonadism (congenital or acquired) in males. Diagnosis of secondary (hypogonadotropic) hypogonadism (congenital or acquired) in males. Hypogonadism (primary or secondary) in males, serum testosterone level. [Man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
Age Restrictions	N/A
Prescriber Restrictions	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of transgender patients.
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. [Note: male is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.] Gender-

PA Criteria	Criteria Details
	Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)-approve. Note: For a patient who has undergone gender reassignment, use this FTM criterion for hypogonadism indication.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Gender-Dysphoric/Gender-Incongruent Persons, Persons Undergoing Female-to-Male (FTM) Gender Reassignment (i.e., Endocrinologic Masculinization)
Part B Prerequisite	No

NUBEQA

Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate cancer - non-metastatic, castration resistant-approve if the requested medication will be used concurrently with a gonadotropin-releasing hormone (GnRH) analog (See Note 1) or if the patient has had a bilateral orchiectomy. Prostate cancer-metastatic, castration sensitive-approve if the medication will be used in combination with a GnRH analog (See Note 1) or if the patient had a bilateral orchiectomy. Note 1: examples of GnRH analogs are leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NUCALA

Products Affected

- NUCALA SUBCUTANEOUS AUTO-
INJECTOR
- NUCALA SUBCUTANEOUS RECON
SOLN
- NUCALA SUBCUTANEOUS SYRINGE
100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another monoclonal antibody therapy.
Required Medical Information	COPD initial - Approve if pt meets all (A, B AND C): A) pt has a blood eosinophil level at least 300 cells per microliter within the previous 6 weeks or prior to treatment with Nucala or another monoclonal antibody therapy that may alter blood eosinophil levels AND B) received at least 3 months of combination therapy with at least two of LAMA, LABA or ICS, AND C) pt meets (i or ii): i) experienced two or more COPD exacerbations requiring treatment with a systemic corticosteroid with or without an antibiotic in the previous 12 months OR ii) experienced one or more COPD exacerbations requiring a hospitalization in the previous 12 months. Cont - Approve if pt meets all (A, B and C): A) has already received at least 6 months of therapy with Nucala AND B) continues to receive combination therapy with an inhaled LABA and LAMA AND C) patient has experienced a beneficial clinical response, defined by one of the following: reduced symptoms, exacerbations, hospitalizations, ED/urgent care visits, improved lung function).
Age Restrictions	(Initial therapy only): Asthma-6 years of age and older. EGPA/Polyps/COPD-18 years of age and older. HES-12 years and older.
Prescriber Restrictions	(Initial therapy only): Asthma/COPD-Prescribed by or in consultation with an allergist, immunologist, or pulmonologist. EGPA-prescribed by or in consultation with an allergist, immunologist, pulmonologist or rheumatologist. HES-prescribed by or in consultation with an allergist, immunologist, hematologist, pulmonologist or rheumatologist. Polyps-prescribed by or in consult with allergist, immunologist or Otolaryngologist
Coverage Duration	Initial-Asthma/polyps/COPD-6 months, EGPA/HES-8 months. 12 months continuation.
Other Criteria	Asthma initial - must have blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 wks (or prior to tx with Nucala or another monoclonal antibody therapy that may lower blood eosinophil levels) AND has received combo tx w/inhaled corticosteroid

PA Criteria	Criteria Details
	<p>AND at least 1 additional asthma controller/maintenance med (Examples: LAMA, LABA, leukotriene receptor antagonist, monoclonal antibody)</p> <p>AND pt's asthma cont to be uncontrolled, or was uncontrolled prior to starting Nucala or another monoclonal antibody therapy for asthma as defined by 1 of following-pt experi 2 or more asthma exacer req tx w/systemic corticosteroids in prev yr, pt experienced 1 or more asthma exacer requiring hospitalization, urgent care visit or ED visit in the prev yr, pt has a FEV1 less than 80 percent predicted, Pt has FEV1/FVC less than 0.80, or Pt's asthma worsens upon taper of oral (systemic) corticosteroid therapy. Cont-pt responded to Nucala tx as determined by the prescribing physician AND Pt cont to receive tx with an inhaled corticosteroid. EGPA initial-approve if pt has active, non-severe disease, has/had a blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 wks or prior to any monoclonal antibody that may lower blood eosinophil levels. Cont-pt responded to Nucala tx as determined by the prescribing physician.HES initial-pt has had hypereosinophilic synd for greater than or equal to 6 months AND has FIP1L1-PDGFRalpha-negative dis AND pt does NOT have identifiable non-hematologic secondary cause of hypereosinophilic synd AND prior to initiating tx with monoclonal antibody that may lower blood eosinophil levels, pt has/had a blood eosinophil level of greater than or equal to 1,000 cells per microliter. Cont-approve if the patient has responded to Nucala tx. Nasal polyps, initial-approve if pt meets ALL of the following criteria(A, B, C and D):A) pt has chronic rhinosinusitis w/nasal polyposis as evidenced by direct examination, endoscopy, or sinus CT scan AND B)pt experienced 2 or more of the following sympt for at least 6 months:nasal congest/obstruct/discharge, and/or reduction/loss of smell AND C)pt meets BOTH of the following (a and b): a)Pt has received tx with intranasal corticosteroid AND b)Pt will continue to receive tx with intranasal corticosteroid concomitantly with Nucala AND D)pt meets 1 of the following (a, b or c): a)Pt has received at least 1 course of tx with a systemic corticosteroid for 5 days or more within the previous 2 years, OR b)Pt has a contraindication to systemic corticosteroid tx, OR c)Pt had prior surgery for nasal polyps.Cont-approve if the pt has received at least 6 months of therapy, continues to receive tx with an intranasal corticosteroid and has responded to tx.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NUPLAZID

Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NURTEC

Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another calcitonin gene-related peptide (CGRP) inhibitor being prescribed for migraine headache prevention if Nurtec ODT is being taken for the preventive treatment of episodic migraine.
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Migraine, Acute treatment (initial and continuation)-approve. Preventive treatment of episodic migraine (initial)-approve if the patient has greater than or equal to 4 but less than 15 migraine headache days per month (prior to initiating a migraine preventive medication). Preventive treatment of episodic migraine (continuation) - approve if the patient has greater than or equal to 4 but less than 15 migraine headache days per month (prior to initiating a migraine preventive medication) and the patient has had a significant clinical benefit from the medication.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NYVEPRIA

Products Affected

- NYVEPRIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Cancer patients receiving chemotherapy-prescribed by or in consultation with an oncologist or hematologist. PBPC-prescribed by or in consultation with an oncologist, hematologist, or physician that specializes in transplantation.
Coverage Duration	Cancer pts receiving chemo-6 mo. PBPC-1 mo
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient meets one of the following: 1) is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), 2) patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia according to the prescribing physician (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status or HIV infection, or 3) patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor and a reduced dose or frequency of chemotherapy may compromise treatment.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients undergoing PBPC collection and therapy
Part B Prerequisite	No

OCTREOTIDE INJECTABLE

Products Affected

- *octreotide acetate injection solution*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Acromegaly-prescr/consult w/endocrinologist. NETs-prescr/consult w/oncologist, endocrinologist, or gastroenterologist. Pheochromocytoma/paraganglioma-prescr/consult w/endo/onc/neuro. Meningioma-prescr/consult w/oncologist, radiologist, neurosurg/thymoma/thymic carcinoma-presc/consult with oncologist. Diarrhea assoc w chemo-presc/consult with oncologist/gastro. Small bowel bleed/angiodyplasia-presc/consult gastroenterologist.
Coverage Duration	Pending CMS Review
Other Criteria	ACROMEGALY (A or B): A) inadequate response to surgery and/or radiotherapy or patient not an appropriate candidate or B) patient is experiencing negative effects due to tumor size (e.g., optic nerve compression) and has pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal based on age and gender. DIARRHEA ASSOC W CHEMO (A and B): A) grade 3 or 4 diarrhea and B) patient has tried at least one antimotility medication. SMALL BOWEL BLEEDS/ANGIODYPLASIA RELATED BLEEDING: pt has chronic, recurrent gastrointestinal bleeds lasting greater than or equal to 3 months.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Meningioma, thymoma and thymic carcinoma, pheochromocytoma and paraganglioma, enterocutaneous fistulas, diarrhea associated with chemotherapy, small bowel bleeds/angiodyplasia related bleeding
Part B Prerequisite	No

ODOMZO

Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	BCC - Must not have had disease progression while on Erivedge (vismodegib).
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Locally advanced BCC approve if the BCC has recurred following surgery/radiation therapy or if the patient is not a candidate for surgery AND the patient is not a candidate for radiation therapy, according to the prescribing physician. Metastatic BCC - approve, if the disease is limited to nodal metastases. (Note-This includes primary or recurrent nodal metastases. A patient with distant metastasis does not meet this requirement.) Diffuse Basal Cell Carcinoma Formation, including basal cell nevus syndrome (Gorlin syndrome) or other genetic forms of multiple basal cell carcinoma - approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Metastatic BCC, diffuse basal cell carcinoma formation
Part B Prerequisite	No

OFEV

Products Affected

- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	IPF-Prescribed by or in consultation with a pulmonologist. Interstitial lung disease associated with systemic sclerosis-prescribed by or in consultation with a pulmonologist or rheumatologist.
Coverage Duration	1 year
Other Criteria	IDIOPATHIC PULMONARY FIBROSIS (IPF), INITIAL [A and B]: A) diagnosis confirmed by presence of usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) or surgical lung biopsy and B) forced vital capacity (FVC) greater than or equal to 40 percent of the predicted value. INTERSTITIAL LUNG DISEASE ASSOCIATED WITH SYSTEMIC SCLEROSIS, INITIAL (A and B): A) diagnosis confirmed by HRCT and B) FVC greater than or equal to 40 percent of the predicted value. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE, INITIAL (all of A, B and C): A) FVC greater than or equal to 45 percent of the predicted value, B) fibrosing lung disease impacting more than 10 percent of lung volume on HRCT, and C) clinical signs of progression. ALL INDICATIONS, CONTINUATION: approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OGSIVEO

Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Desmoid tumors (aggressive fibromatosis)-approve if the patient has progressing desmoid tumors, the desmoid tumors are not amenable to surgery or radiotherapy and if the patient requires systemic treatment. Note: Progressing desmoid tumors are defined as greater than or equal to 20 percent progression within 12 months
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OJEMDA

Products Affected

- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)
- OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	6 months of age and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	PEDIATRIC LOW GRADE GLIOMA-patient has relapsed or refractory disease and the tumor is positive for one of the following: BRAF fusion, BRAF rearrangement or BRAF V600 mutation.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OJJAARA

Products Affected

- OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Myelofibrosis-approve if the patient has (A, B or C): A) intermediate-risk or high-risk disease, or B) lower-risk disease and has one disease-related symptom (Examples: fatigue, fever, night sweats, weight loss, abdominal discomfort, splenomegaly, thrombocytosis, or leukocytosis), or C) myelofibrosis-associated anemia. Accelerated or blast phase myeloproliferative neoplasm- approve if the patient has at least one disease- related symptom (Examples: fatigue, fever, night sweats, weight loss, abdominal discomfort, splenomegaly, thrombocytosis, or leukocytosis).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Accelerated or blast phase myeloproliferative neoplasm
Part B Prerequisite	No

ONUREG

Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML - Approve if the medication is used for post-remission maintenance therapy AND allogeneic hematopoietic stem cell transplant is not planned. Peripheral T-cell lymphoma - all of (A, B, and C): A) relapsed or refractory disease, and B) pt has one of the following (i, ii or iii): i) angioimmunoblastic T-cell lymphoma, or ii) nodal peripheral T-cell lymphoma with T-follicular helper (TFH) phenotype, or iii) follicular T-cell lymphoma, and C) medication is used as a single agent.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Peripheral T-cell lymphoma
Part B Prerequisite	No

OPSUMIT

Products Affected

- OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	PAH WHO group, right heart catheterization results
Age Restrictions	N/A
Prescriber Restrictions	PAH - must be prescribed by or in consultation with a cardiologist or a pulmonologist.
Coverage Duration	Authorization will be for 1 year
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1 patients are required to have had a right-heart catheterization to confirm the diagnosis of PAH to ensure appropriate medical assessment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OPSYNVI

Products Affected

- OPSYNVI

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with guanylate cyclase stimulators
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a pulmonologist (initial/continuation)
Coverage Duration	1 year
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1-approve if patient has had a right-heart catheterization to confirm the diagnosis.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORENCIA

Products Affected

- ORENCEA CLICKJECT
- ORENCEA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	N/A
Prescriber Restrictions	Initial therapy only-RA and JIA/JRA prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or dermatologist.
Coverage Duration	Approve through end of plan year
Other Criteria	RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). PsA-approve. Juvenile idiopathic arthritis (JIA) [or Juvenile Rheumatoid Arthritis (JRA)]-approve if the patient has tried one other agent for this condition or the patient will be starting on Orencia concurrently with methotrexate, sulfasalazine or leflunomide or the patient has an absolute contraindication to methotrexate, sulfasalazine or leflunomide or the patient has aggressive disease. Cont tx - responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORGOVYX

Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prostate Cancer-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORKAMBI

Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Kalydeco, Trikafta or Symdeko.
Required Medical Information	N/A
Age Restrictions	1 year of age and older
Prescriber Restrictions	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	1 year
Other Criteria	CF - Approve if the pt meets A, B and C: A) pt has two copies of the F508del mutation in the CFTR gene, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORSERDU

Products Affected

- ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast cancer in postmenopausal women or Men-approve if the patient meets the following criteria (A, B, C, D, and E): A) Patient has recurrent or metastatic disease, AND B) Patient has estrogen receptor positive (ER+) disease, AND C) Patient has human epidermal growth factor receptor 2 (HER2)-negative disease, AND D) Patient has estrogen receptor 1 gene (ESR1)-mutated disease, AND E) Patient has tried at least one endocrine therapy. Note: Examples of endocrine therapy include fulvestrant, anastrozole, exemestane, letrozole, and tamoxifen.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OTEZLA

Products Affected

- OTEZLA MG (51), 10 MG (4)-20 MG (4)-30 MG (47)
- OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with a Biologic or with a Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARD).
Required Medical Information	Diagnosis, previous drugs tried
Age Restrictions	PP/PsA- 6 years and older (initial), All other dx - 18 years and older (initial)
Prescriber Restrictions	All dx-initial only-PsA - Prescribed by or in consultation with a dermatologist or rheumatologist. PP - prescribed by or in consultation with a dermatologist. Behcet's-prescribed by or in consultation with a dermatologist or rheumatologist
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: PLAQUE PSORIASIS (PP) [A, B or C]: A) tried at least one traditional systemic agent (e.g., methotrexate [MTX], cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial with a biologic also counts) or B) contraindication to MTX or C) patient has mild to moderate disease and the patient requires systemic therapy. PSORIATIC ARTHRITIS (PsA): approve. BEHCET'S-oral ulcers or other mucocutaneous involvement. CONTINUATION THERAPY (PP/PsA/Behcet's): received 4 months of therapy and had a response.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OXERVATE

Products Affected

- OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 16 weeks per affected eye(s)
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by an ophthalmologist or an optometrist.
Coverage Duration	Initial-8 weeks, continuation-approve for an additional 8 weeks
Other Criteria	Patients who have already received Oxervate-approve if the patient has previously received less than or equal to 8 weeks of treatment per affected eye(s) and the patient has a recurrence of neurotrophic keratitis.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PANRETIN

Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist, oncologist, or infectious disease specialist
Coverage Duration	1 year
Other Criteria	Kaposi Sarcoma-approve if the patient is not receiving systemic therapy for Kaposi Sarcoma.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PEMAZYRE

Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Cholangiocarcinoma-approve if the patient has unresectable locally advanced or metastatic disease and the tumor has a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement, as detected by an approved test AND the patient has been previously treated with at least one systemic therapy regimen. Myeloid/lymphoid neoplasms-approve if the patient has eosinophilia and the cancer has fibroblast growth factor receptor 1 (FGFR1) rearrangement, as detected by an approved test and the cancer is in chronic phase or blast phase.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PENICILLAMINE

Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Wilson's Disease-Prescribed by or in consultation with a gastroenterologist, hepatologist or liver transplant physician
Coverage Duration	1 year
Other Criteria	Cystinuria-approve. Wilson's disease-approve if diagnosis is confirmed by ONE of the following (i or ii): i. Genetic testing results confirming biallelic pathogenic ATP7B mutations (in either symptomatic or asymptomatic individuals), OR ii. Confirmation of at least two of the following (a, b, c, d): a) Presence of Kayser-Fleischer rings, b) Serum ceruloplasmin level less than 20 mg/dL, c) Liver biopsy findings consistent with Wilson's disease, d) 24-hour urinary copper greater than 40 mcg/24 hours.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PHENYLBUTYRATE

Products Affected

- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant therapy with more than one phenylbutyrate product
Required Medical Information	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a metabolic disease specialist (or specialist who focuses in the treatment of metabolic diseases)
Coverage Duration	Pt meets criteria with no genetic test - 3 mo approval. Pt had genetic test - 12 mo approval
Other Criteria	Urea cycle disorders-approve if genetic or enzymatic testing confirmed a urea cycle disorder or if the patient has hyperammonemia diagnosed with an ammonia level above the upper limit of the normal reference range for the reporting laboratory.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PHEOCHROMOCYTOMA

Products Affected

- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior medication trials
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the management of pheochromocytoma (initial and continuation therapy for metyrosine)
Coverage Duration	Authorization will be for 1 year
Other Criteria	If the requested drug is metyrosine for initial therapy, approve if the patient has tried a selective alpha blocker (e.g., doxazosin, terazosin or prazosin). If the requested drug is metyrosine for continuation therapy, approve if the patient is currently receiving metyrosine or has received metyrosine in the past.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PHOSPHODIESTERASE-5 INHIBITORS FOR PAH

Products Affected

- *alyq*
- *sildenafil (pulmonary arterial hypertension) oral tablet*
- *tadalafil (pulmonary arterial hypertension) oral tablet 20 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use With Guanylate Cyclase Stimulators.
Required Medical Information	Diagnosis, right heart cath results
Age Restrictions	N/A
Prescriber Restrictions	For PAH, if prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Clinical criteria incorporated into the quantity limit edits for sildenafil 20 mg tablets require confirmation that the indication is PAH (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PIQRAY

Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 300 MG/DAY (150 MG X 2)
- 250 MG/DAY (200 MG X 1)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast Cancer. Approve if the patient meets the following criteria (A, B, C, D, E and F): A) The patient is a postmenopausal female, male or pre/perimenopausal and is receiving ovarian suppression with a gonadotropin-releasing hormone (GnRH) agonist or has had surgical bilateral oophorectomy or ovarian irradiation AND B) The patient has advanced or metastatic hormone receptor (HR)-positive disease AND C) The patient has human epidermal growth factor receptor 2 (HER2)-negative disease AND D) The patient has PIK3CA-mutated breast cancer as detected by an approved test AND E) The patient has progressed on or after at least one prior endocrine-based regimen (e.g., anastrozole, letrozole, exemestane, tamoxifen, toremifene or fulvestrant) AND F) Piqray will be used in combination with fulvestrant injection.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PIRFENIDONE

Products Affected

- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 801 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	1 year
Other Criteria	IPF (initial therapy)- must have FVC greater than or equal to 40 percent of the predicted value AND IPF must be diagnosed with either findings on high-resolution computed tomography (HRCT) indicating usual interstitial pneumonia (UIP) or surgical lung biopsy demonstrating UIP. IPF (continuation of therapy)-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PLEGRIDY

Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of with other disease-modifying agents used for multiple sclerosis (MS).
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by, or in consultation with, a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

POMALYST

Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Kaposi Sarcoma/MM-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	CNS Lymphoma-approve if the patient has relapsed or refractory disease. Kaposi Sarcoma-Approve if the patient meets one of the following (i or ii): i. patient is Human Immunodeficiency Virus (HIV)-negative OR ii. patient meets both of the following (a and b): a) The patient is Human Immunodeficiency Virus (HIV)-positive AND b) The patient continues to receive highly active antiretroviral therapy (HAART). MM-approve if the patient has received at least one other Revlimid (lenalidomide tablets)-containing regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Systemic Light Chain Amyloidosis, Central Nervous System (CNS) Lymphoma
Part B Prerequisite	No

POSACONAZOLE (ORAL)

Products Affected

- *posaconazole oral tablet, delayed release (dr/ec)*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Aspergillus/Candida proph, mucormycosis-6 mo, all others-3 months
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	mucomycosis - maintenance, fusariosis, invasive - treatment fungal infections (systemic) in patients with human immunodeficiency virus (HIV) infections (e.g., histoplasmosis, coccidioidomycosis) - treatment.
Part B Prerequisite	No

PREVYMIS

Products Affected

- PREVYMIS ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PROLIA

Products Affected

- CONEXXENCE
- PROLIA
- JUBBONTI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other medications for osteoporosis
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Treatment of postmenopausal osteoporosis/Treatment of osteoporosis in men (to increase bone mass) [a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression], approve if the patient meets one of the following: 1. has had inadequate response after 12 months of therapy with an oral bisphosphonate, had osteoporotic fracture or fragility fracture while receiving an oral bisphosphonate, or intolerance to an oral bisphosphonate, OR 2. the patient cannot take an oral bisphosphonate because they cannot swallow or have difficulty swallowing, they cannot remain in an upright position, or they have a pre-existing GI medical condition, OR 3. pt has tried an IV bisphosphonate (ibandronate or zoledronic acid), OR 4. the patient has severe renal impairment (eg, creatinine clearance less than 35 mL/min) or chronic kidney disease, or if the patient has an osteoporotic fracture or fragility fracture . Treatment of bone loss in patient at high risk for fracture receiving ADT for nonmetastatic prostate cancer, approve if the patient has prostate cancer that is not metastatic to the bone and the patient is receiving ADT (eg, leuprolide, triptorelin, goserelin) or the patient has undergone a bilateral orchiectomy. Treatment of bone loss (to increase bone mass) in patients at high risk for fracture receiving adjuvant AI therapy for breast cancer, approve if the patient has breast cancer that is not metastatic to the bone and in receiving concurrent AI therapy (eg, anastrozole, letrozole,

PA Criteria	Criteria Details
	exemestane). Treatment of GIO, approve if pt tried one oral bisphosphonate OR pt cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing or the patient cannot remain in an upright position post oral bisphosphonate administration or has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR pt has tried zoledronic acid (Reclast), OR pt has severe renal impairment (CrCL less than 35 mL/min) or has CKD or has had an osteoporotic fracture or fragility fracture.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PROMACTA

Products Affected

- *eltrombopag olamine*
- PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Cause of thrombocytopenia. Thrombocytopenia due to HCV-related cirrhosis, platelet counts. Severe aplastic anemia, platelet counts and prior therapy. MDS-platelet counts.
Age Restrictions	N/A
Prescriber Restrictions	Immune Thrombocytopenia or Aplastic Anemia, prescribed by, or after consultation with, a hematologist (initial therapy). Hep C, prescribed by, or after consultation with, a gastroenterologist, hematologist, hepatologist, or a physician who specializes in infectious disease (initial therapy). MDS-presc or after consult with heme/onc (initial therapy). Post-transplant, prescribed by or in consult with a hematologist, oncologist or stem cell transplant specialist physician (initial)
Coverage Duration	Immune Thrombo/MDS init3mo,cont1yr,AAinit4mo,cont1yr,Thrombo/HepC1yr,Transplant-init3mo,cont6mo
Other Criteria	Thrombocytopenia in patients with immune thrombocytopenia, initial-approve if the patient has a platelet count less than 30, 000 microliters or less than 50, 000 microliters and the patient is at an increased risk for bleeding AND the patient has tried ONE other therapy (e.g., systemic corticosteroids, intravenous immunoglobulin, anti-D immunoglobulin, Nplate, Tavalisse, Doptelet, rituximab) or has undergone a splenectomy. Cont-approve if the patient demonstrates a beneficial clinical response and remains at risk for bleeding complications. Treatment of thrombocytopenia in patients with Chronic Hepatitis C initial - approve if the patient will be receiving interferon-based therapy for chronic hepatitis C AND if the patient has low platelet counts at baseline (eg, less than 75,000 microliters). Aplastic anemia initial - approve if the patient has low platelet counts at baseline/pretreatment (e.g., less than 30,000 microliter) AND tried one immunosuppressant therapy (e.g., cyclosporine) OR patient will be using Promacta in combination with standard immunosuppressive therapy. Cont-approve if the patient demonstrates a beneficial clinical response. MDS

PA Criteria	Criteria Details
	initial-approve if patient has low- to intermediate-risk MDS AND the patient has a platelet count less than 30, 000 microliters or less than 50, 000 microliters and is at an increased risk for bleeding. Cont-approve if the patient demonstrates a beneficial clinical response and remains at risk for bleeding complications. Thrombocytopenia post-allogeneic transplantation, initial - approve if, according to the prescriber, the patient has poor graft function AND has a platelet count less than 50,000/mcL. Cont- patient demonstrated a beneficial clinical response.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Thrombocytopenia in Myelodysplastic Syndrome (MDS), Thrombocytopenia in a patient post-allogeneic transplantation
Part B Prerequisite	No

PYRIMETHAMINE

Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient's immune status (Toxoplasma gondii Encephalitis, chronic maintenance and prophylaxis, primary)
Age Restrictions	N/A
Prescriber Restrictions	Toxoplasma gondii Encephalitis, Chronic Maintenance and Prophylaxis (Primary)-prescribed by or in consultation with an infectious diseases specialist. Toxoplasmosis Treatment-prescribed by or in consultation with an infectious diseases specialist, a maternal-fetal medicine specialist, or an ophthalmologist.
Coverage Duration	12 months
Other Criteria	Toxoplasma gondii Encephalitis, Chronic Maintenance, approve if the patient is immunosuppressed. Toxoplasma gondii Encephalitis Prophylaxis (Primary), approve if the patient is immunosuppressed.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chronic maintenance and prophylaxis of Toxoplasma Gondii encephalitis
Part B Prerequisite	No

QINLOCK

Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Gastrointestinal stromal tumor (GIST)-approve if the patient has tried imatinib or avapritinib tablets, AND the patient meets one of the following criteria (i, ii, or iii): i. Patient has tried sunitinib and regorafenib tablets, OR ii. Patient has tried dasatinib tablets, OR iii. Patient is intolerant of sunitinib. Melanoma, cutaneous-approve if the patient has metastatic or unresectable disease, AND the patient has an activating KIT mutation, AND the patient has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Melanoma, cutaneous
Part B Prerequisite	No

QULIPTA

Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another calcitonin gene-related peptide (CGRP) inhibitor being prescribed for migraine headache prevention
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Migraine headache prevention-approve if the patient meets (A and B): A) has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication, and B) if the pt is currently taking Qulipta, the pt has had significant clinical benefit.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RADICAVA ORS

Products Affected

- RADICAVA ORS STARTER KIT SUSP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	ALSFRS-R score
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, a neuromuscular disease specialist, or a physician specializing in the treatment of ALS (initial and continuation).
Coverage Duration	Initial, 6 months. Continuation, 6 months
Other Criteria	ALS, initial therapy - approve if the patient meets ALL of the following criteria: 1. According to the prescribing physician, the patient has a definite or probable diagnosis of ALS, based on the application of the El Escorial or the revised Airlie house diagnostic criteria AND 2. Patient has a score of two points or more on each item of the ALS Functional Rating Scale - Revised (ALSFRS-R) [ie, has retained most or all activities of daily living], AND 3. Patient has adequate respiratory function according to the prescriber, AND 4. Patient has received or is currently receiving riluzole tablets. Note-a trial of Tiglutik or Exservan would also count. ALS, continuation therapy - approve if, according to the prescribing physician, the patient continues to benefit from therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RELEUKO

Products Affected

- RELEUKO SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Cancer/AML, MDS, ALL-oncologist or a hematologist. Cancer patients receiving BMT and PBPC-prescribed by or in consultation with an oncologist, hematologist, or a physician who specializes in transplantation. SCN-hematologist. HIV/AIDS neutropenia-infectious disease (ID) physician (MD), hematologist, or MD specializing in HIV/AIDS. Radiation syndrome-prescribed by or in consultation with expert in acute radiation.
Coverage Duration	chemo/SCN/AML-6 mo.HIV/AIDS-4 mo.MDS-3 mo.PBPC,Drug induce A/N, ALL,BMT,Radi-1 mo
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient meets one of the following conditions: 1)patient is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20 percent based on the chemotherapy regimen), 2)patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20 percent based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia (eg, aged greater than or equal to 65 years, priorchemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status, or HIV infection), 3)patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor (eg, Leukine, filgrastim products, pegfilgrastim products and a reduced dose or frequency of chemotherapy may compromise treatment, or 4)patient has received chemotherapy, has febrile neutropenia and has at least one risk factor (eg, sepsis syndrome, aged greater than 65 years, severe neutropenia

PA Criteria	Criteria Details
	[absolute neutrophil count less than 100 cells/mm ³], neutropenia expected to be greater than 10 days in duration, invasive fungal infection)
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Neutropenia associated with HIV or AIDS, Treatment of myelodysplastic syndromes (MDS), Drug induced agranulocytosis or neutropenia, Acute lymphocytic leukemia (ALL), Radiation Syndrome (Hematopoietic Syndrome of Acute Radiation Syndrome), peripheral blood progenitor cell transplantation in patients with cancer
Part B Prerequisite	No

REPATHA

Products Affected

- REPATHA
- REPATHA SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of Leqvio or Praluent.
Required Medical Information	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history (as specified in the Other Criteria field)
Age Restrictions	ASCVD/Primary Hyperlipidemia - 18 yo and older, HoFH/HeFH - 10 yo and older.
Prescriber Restrictions	N/A
Coverage Duration	Approve for 1 year
Other Criteria	Pending CMS Review
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RETEVMO

Products Affected

- RETEVMO ORAL TABLET 120 MG,
160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Medullary Thyroid Cancer/Thyroid Cancer/Solid tumors-2 years and older, all others 18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-Approve if the patient has recurrent, advanced or metastatic disease AND the tumor is RET fusion-positive. Thyroid cancer-approve if the patient has rearranged during transfection (RET) fusion positive or RET mutation positive disease or RET pathogenic variant AND the patient meets i, ii or iii: i. patient has anaplastic thyroid cancer OR ii. the disease requires treatment with systemic therapy and patient has medullary thyroid cancer or iii. the disease requires treatment with systemic therapy and the disease is radioactive iodine-refractory if radioactive iodine is appropriate. Solid tumors-approve if the patient has recurrent, advanced or metastatic disease and the tumor is rearranged during transfection (RET) fusion-positive. Histiocytic neoplasm-approve if the patient has a rearranged during transfection (RET) fusion and has Langerhans cell histiocytosis or Erdheim Chester disease or Rosai-Dorfman disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Anaplastic thyroid carcinoma, histiocytic neoplasm
Part B Prerequisite	No

REVUFORJ

Products Affected

- REVUFORJ ORAL TABLET 110 MG,
160 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	1 year and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	ACUTE LEUKEMIA-patient has relapsed or refractory disease and the disease is positive for a lysine methyltransferase 2A (KMT2A) gene translocation.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

REZDIFFRA

Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (initial)
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist, gastroenterologist, or hepatologist (initial/continuation)
Coverage Duration	1 year
Other Criteria	<p>INITIAL THERAPY: METABOLIC-DYSFUNCTION ASSOCIATED STEATOHEPATITIS (MASH)/NON-ALCOHOLIC STEATOHEPATITIS (NASH) WITH MODERATE-ADVANCED LIVER FIBROSIS: All of (i, ii and iii): i) Diagnosed by (a or b): a) Liver biopsy performed within 3 years preceding treatment with Rezdiffra showing non-alcoholic fatty liver disease activity score of greater than or equal to 4 with a score of greater than or equal to 1 in ALL of the following: steatosis, ballooning and lobular inflammation, or b) One of the following within 6 months preceding treatment with Rezdiffra (1, 2 or 3): 1) Elastography (e.g. vibration-controlled transient elastography (e.g., FibroScan), transient elastography, magnetic resonance elastography, acoustic radiation force impulse imaging, shear wave elastography) or 2) Computed tomography or 3) Magnetic resonance imaging, and ii) stage F2 or F3 fibrosis prior to Rezdiffra and iii) This will be used in combination with appropriate diet and exercise therapy (prescriber attestation the patient has received counseling on diet and exercise). CONTINUATION THERAPY (on therapy less than 1 year or restarting, review as initial therapy): MASH/NASH: All of (i, ii and iii): i) completed greater than or equal to 1 year of therapy and has not had worsening of fibrosis or MASH/NASH, and ii) has not progressed to stage F4 (cirrhosis) and iii) This will be used in combination with appropriate diet and exercise therapy (prescriber attestation the patient has received counseling on diet and exercise).</p>

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

REZLIDHIA

Products Affected

- REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Acute myeloid leukemia-approve if the patient has relapsed or refractory disease and the patient has isocitrate dehydrogenase-1 (IDH1) mutation positive disease as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

REZUROCK

Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	12 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Graft-versus-host disease-approve if the patient has chronic graft-versus-host disease and has tried at least two conventional systemic treatments (e.g., ibrutinib, cyclosporine) for chronic graft-versus-host disease.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RILUZOLE

Products Affected

- *riluzole*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, a neuromuscular disease specialist, or a physician specializing in the treatment of ALS.
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RINVOQ

Products Affected

- RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a targeted synthetic DMARD, Concurrent use with other potent immunosuppressants, Concurrent use with an anti-interleukin monoclonal antibody, Concurrent use with other janus kinase inhibitors, or concurrent use with a biologic immunomodulator.
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	PsA/JIA - 2 years and older (initial therapy), RA/UC/AS/CD/nr-axSpA/GCA-18 years and older (initial therapy), AD-12 years and older (initial therapy)
Prescriber Restrictions	RA/AS/Non-Radiographic Spondy/JIA/GCA, prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or a dermatologist. AD-prescr/consult with allergist, immunologist or derm. UC/CD-prescribed by or in consultation with a gastroenterologist. (all apply to initial therapy only)
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: RHEUMATOID ARTHRITIS (RA)/PSORIATIC ARTHRITIS (PsA)/ULCERATIVE COLITIS (UC)/ANKYLOSING SPONDYLITIS (AS)/CROHN'S DISEASE (CD)/ JUVENILE IDIOPATHIC ARTHRITIS (JIA): 3-month trial of at least one tumor necrosis factor inhibitor (TNFi) or unable to tolerate a 3-month trial. ATOPIC DERMATITIS (AD): 90 day trial of at least one systemic therapy (e.g., Dupixent [dupilumab subcutaneous injection] and Adbry [tralokinumab-ldrm subcutaneous injection]. Azathioprine, cyclosporine, or mycophenolate mofetil also count.) or unable to tolerate a 90 day trial. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (A and B): A) objective signs of inflammation defined as having at least one of the following (a or b): a) C-reactive protein elevated beyond the upper limit of normal or b) sacroiliitis reported on MRI and B) 3-month trial of at least

PA Criteria	Criteria Details
	one TNFi or was unable to tolerate a 3-month trial. GIANT CELL ARTERITIS: tried or is currently taking one systemic corticosteroid or corticosteroids are contraindicated. CONTINUATION THERAPY: ALL INDICATIONS: patient responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RINVOQ LQ

Products Affected

- RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a targeted synthetic DMARD, other potent immunosuppressants, other janus kinase inhibitors, or a biologic immunomodulator.
Required Medical Information	Diagnosis
Age Restrictions	PsA-2 years and older (initial therapy)
Prescriber Restrictions	JIA-prescribed by or in consultation with a rheumatologist (initial therapy). PsA-prescribed by or in consultation with a rheumatologist or a dermatologist (initial therapy)
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: JUVENILE IDIOPATHIC ARTHRITIS (JIA)/ PSORIATIC ARTHRITIS (PsA) - 3-month trial of at least one tumor necrosis factor inhibitor (TNFi) or unable to tolerate a 3-month trial. CONTINUATION THERAPY: ALL INDICATIONS - patient responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ROFLUMILAST (ORAL)

Products Affected

- *roflumilast*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD), medications tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year.
Other Criteria	COPD, approve in patients who meet all of the following conditions: Patients has severe COPD or very severe COPD, AND Patient has a history of exacerbations, AND Patient has tried a medication from two of the three following drug categories: long-acting beta2-agonist (LABA) [eg, salmeterol, indacaterol], long-acting muscarinic antagonist (LAMA) [eg, tiotropium], inhaled corticosteroid (eg, fluticasone).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ROMVIMZA

Products Affected

- ROMVIMZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	TENOSYNOVIAL GIANT CELL TUMOR (PIGMENTED VILLONODULAR SYNOVITIS)-tumor is not amenable to improvement with surgery.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ROZLYTREK

Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	NSCLC-18 years and older, Solid Tumors-1 month and older, Pediatric Diffuse High-Grade Glioma-less than 18 years old
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Solid Tumors-Approve if the patient's tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the tumor is metastatic OR surgical resection of tumor will likely result in severe morbidity. Non-Small Cell Lung Cancer-Approve if the patient has ROS1-positive metastatic disease and the mutation was detected by an approved test. Pediatric Diffuse High-Grade Glioma- approve if the tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the medication is used either as adjuvant therapy or for recurrent or progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pediatric Diffuse High-Grade Glioma
Part B Prerequisite	No

RUBRACA

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Rubraca is being used. BRCA-mutation (germline or somatic) status. Other medications tried for the diagnosis provided
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Maintenance Therapy of Ovarian, Fallopian tube or Primary peritoneal cancer-Approve if the patient is in complete or partial response after a platinum-based chemotherapy regimen and the patient is in complete or partial response to first-line primary treatment or if the patient has recurrent disease and has a BRCA mutation. Castration-Resistant Prostate Cancer - Approve if the patient meets the following criteria (A, B and C): A) The patient has metastatic disease that is BRCA-mutation positive (germline and/or somatic) AND B) The patient meets one of the following criteria (i or ii): i. The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog OR ii. The patient has had a bilateral orchiectomy AND C) The patient has been previously treated with at least one androgen receptor-directed therapy. Pancreatic adenocarcinoma-approve if pt has a BRCA mutation or PALB2 mutation AND pt has tried platinum-based chemotherapy AND has not had disease progression following the most recent platinum-based chemotherapy. Uterine leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Uterine Leiomyosarcoma, Pancreatic Adenocarcinoma
Part B Prerequisite	No

RUFINAMIDE

Products Affected

- *rufinamide*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Patients 1 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Initial therapy-approve if rufinamide is being used for adjunctive treatment. Continuation-approve if the patient is responding to therapy
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Treatment-Refactory Seizures/Epilepsy
Part B Prerequisite	No

RYDAPT

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For AML, FLT3 status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML -approve if the patient is FLT3-mutation positive as detected by an approved test. Myeloid or lymphoid Neoplasms with eosinophilia-approve if the patient has an FGFR1 rearrangement or has an FLT3 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myeloid or lymphoid Neoplasms with eosinophilia
Part B Prerequisite	No

SAPROPTERIN

Products Affected

- *sapropterin*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Palynziq
Required Medical Information	Diagnosis, Phe concentration
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a specialist who focuses in the treatment of metabolic diseases (initial therapy)
Coverage Duration	Initial-12 weeks, Continuation-1 year
Other Criteria	Initial - approve. Continuation (Note-if the patient has received less than 12 weeks of therapy or is restarting therapy with sapropterin should be reviewed under initial therapy) - approve if the patient has had a response to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SCEMBLIX

Products Affected

- SCEMBLIX ORAL TABLET 100 MG,
20 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Chronic Myeloid Leukemia (CML)-approve if the patient meets the following (A and B): A) Patient has Philadelphia chromosome-positive or BCR::ABL1-positive chronic myeloid leukemia, AND B) Patient meets one of the following (i, ii or iii): i. Patient has newly diagnosed disease, OR ii. The chronic myeloid leukemia is T315I-positive, OR iii. Patient has tried at least one other tyrosine kinase inhibitor. Note: Examples of tyrosine kinase inhibitors include imatinib, Bosulif (bosutinib tablets), Iclusig (ponatinib tablets), dasatinib, and nilotinib capsules. Myeloid/Lymphoid Neoplasms with Eosinophilia - approve if the tumor has an ABL1 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myeloid/Lymphoid Neoplasms with Eosinophilia
Part B Prerequisite	No

SIGNIFOR

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician or specializes in the treatment of Cushing's syndrome (initial therapy)
Coverage Duration	Cushing's disease/syndrome-Initial therapy - 4 months, Continuation therapy - 1 year.
Other Criteria	Cushing's disease, initial therapy - approve if, according to the prescribing physician, the patient is not a candidate for surgery, or surgery has not been curative. Cushing's disease, continuation therapy - approve if the patient has already been started on Signifor/Signifor LAR and, according to the prescribing physician, the patient has had a response and continuation of therapy is needed to maintain response.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SIRTURO

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	Patients weighing less than 8 kg
Required Medical Information	Diagnosis, concomitant therapy
Age Restrictions	Patients 2 years of age or older
Prescriber Restrictions	Pending CMS Review
Coverage Duration	9 months
Other Criteria	Tuberculosis (Pulmonary) -Approve if the patient has multidrug-resistant tuberculosis or Mycobacterium tuberculosis resistant to at least rifampin and isoniazid, and the requested medication is prescribed as part of a combination regimen with other anti-tuberculosis agents
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SKYRIZI

Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE
- SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
Required Medical Information	Diagnosis, Previous medication use
Age Restrictions	PP/UC-18 years of age and older (initial therapy)
Prescriber Restrictions	PP-Prescribed by or in consultation with a dermatologist (initial therapy), PsA-prescribed by or in consultation with a rheumatologist or dermatologist (initial therapy), CD/UC-presc/consult-gastro (initial therapy)
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent for psoriasis (e.g., methotrexate [MTX], cyclosporine, acitretin tablets, or PUVA) for at least 3 months, unless intolerant. (Note: a 3-month trial or previous intolerance to at least one biologic also counts) or B) contraindication to MTX. PSORIATIC ARTHRITIS (PsA): approve. CROHN'S DISEASE (CD):approve. ULCERATIVE COLITIS (UC)-approve. CD/UC: Patient must be receiving induction dosing with Skyrizi IV within 3 months of initiating therapy with Skyrizi subcutaneous. CONTINUATION THERAPY: ALL INDICATIONS: patient has responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR

Products Affected

- SOFOSBUVIR-VELPATASVIR

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with other direct acting antivirals, excluding ribavirin.
Required Medical Information	Diagnosis
Age Restrictions	3 years or older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
Coverage Duration	Criteria will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	Criteria will be applied according to AASLD guidelines.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance
Part B Prerequisite	No

SOLARAZE

Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 6 months.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SOMAVERT

Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapy, concomitant therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	1 year
Other Criteria	ACROMEGALY (A or B): A) inadequate response to surgery and/or radiotherapy or patient not an appropriate candidate or B) patient is experiencing negative effects due to tumor size (e.g., optic nerve compression) and has pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal based on age and gender.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SORAFENIB

Products Affected

- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Bone cancer, approve if the patient has recurrent chordoma or has osteosarcoma and has tried one standard chemotherapy regimen. GIST, approve if the patient has tried TWO of the following: imatinib mesylate, avapritinib, sunitinib, dasatinib, ripretinib or regorafenib. Differentiated (ie, papillary, follicular, oncocytic) thyroid carcinoma (DTC), approve if the patient is refractory to radioactive iodine treatment. Medullary thyroid carcinoma, approve if the patient has tried at least one systemic therapy. AML - Approve if disease is FLT3-ITD mutation positive as detected by an approved test and the medication is used in combination with azacitidine or decitabine or patient has had an allogeneic stem cell transplant and is in remission. Renal cell carcinoma (RCC)-approve if the patient has relapsed or advanced clear cell histology and the patient has tried at least one systemic therapy (e.g., Inlyta, Votrient, Sutent Cabometyx). Ovarian, fallopian tube, primary peritoneal cancer-approve if the patient has platinum resistant disease and sorafenib is used in combination with topotecan. HCC-approve if the patient has unresectable or metastatic disease. Soft tissue sarcoma-approve if the patient has angiosarcoma or desmoid tumors (aggressive fibromatosis) or solitary fibrous tumor/hemangiopericytoma. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an FLT3 rearrangement. Please note for all diagnoses: Part B before Part D Step Therapy applies only to beneficiaries enrolled in an MA-PD plan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Bone cancer, Soft tissue sarcoma, gastrointestinal stromal tumors (GIST), medullary thyroid carcinoma, Acute Myeloid Leukemia, ovarian, fallopian tube, primary peritoneal cancer, myeloid/lymphoid neoplasms with eosinophilia
Part B Prerequisite	Yes

SOTYKTU

Products Affected

- SOTYKTU

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other biologics or with targeted synthetic disease-modifying antirheumatic drugs (DMARDs). Concurrent use with other potent immunosuppressants, including methotrexate.
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (initial)
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist (initial)
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY: PLAQUE PSORIASIS (PP) [A or B]: A) tried at least one traditional systemic agent (e.g., MTX, cyclosporine (CSA), acitretin, PUVA) for at least 3 months, unless intolerant (Note: a trial of a biologic will also count) or B) contraindication to MTX. CONTINUATION THERAPY: patient had a response to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SPRYCEL

Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*
- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which dasatinib is being used. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For melanoma, cutaneous- KIT mutation and previous therapies.
Age Restrictions	GIST/bone cancer/ melanoma, cutaneous-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	For CML, patient must have Ph-positive or BCR::ABL1-positive CML. For ALL, patient must have Ph-positive ALL or ABL-class translocation. For Bone Cancer-approve if patient has chondrosarcoma or chordoma. GIST - approve if the patient has tried imatinib or avapritinib. For melanoma, cutaneous - approve if patient has metastatic or unresectable disease AND has an activating KIT mutation AND has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	GIST, bone cancer, melanoma cutaneous
Part B Prerequisite	No

STELARA

Products Affected

- SELARSDI SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML
- STELARA SUBCUTANEOUS SOLUTION
- STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML
- YESINTEK SUBCUTANEOUS SOLUTION
- YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	Ustekinumab should not be given in combination with a Biologic DMARD or Targeted Synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	PP-6 years and older (initial therapy).
Prescriber Restrictions	Plaque psoriasis. Prescribed by or in consultation with a dermatologist (initial therapy). PsA-prescribed by or in consultation with a rheumatologist or dermatologist (initial therapy). CD/UC-prescribed by or in consultation with a gastroenterologist (initial therapy).
Coverage Duration	Approve through end of plan year
Other Criteria	INITIAL THERAPY for USTEKINUMAB SC: PLAQUE PSORIASIS (PP) [A or B]: A) tried one traditional systemic agent (e.g., methotrexate [MTX], cyclosporine, acitretin, psoralen plus PUVA) for at least 3 months, unless intolerant or B) contraindication to MTX. (Note: a 3-month trial or intolerance of at least one biologic that is not Stelara or a Stelara biosimilar also counts.) CROHN'S DISEASE (CD): approve if pt receiving/received single IV loading dose within 2 months of initiating therapy with ustekinumab SC. ULCERATIVE COLITIS (UC): receiving/received single IV loading dose within 2 months of initiating therapy with ustekinumab SC. CONTINUATION THERAPY: PP/PsA/CD/UC: patient has responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

STIVARGA

Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Stivarga is being used. Prior therapies tried. For metastatic CRC, KRAS/NRAS mutation status.
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For GIST, (A or B): A) patient has previously been treated with (i and ii): i) imatinib or Ayvakit and ii) sunitinib or Sprycel, or B) medication is used as first-line therapy for succinate dehydrogenase (SDH)-deficient disease. For HCC, patient must have previously been treated with at least one systemic regimen. Soft tissue sarcoma, advanced or metastatic disease-approve if the patient has non-adipocytic sarcoma, angiosarcoma, or pleomorphic rhabdomyosarcoma. Bone Cancer-approve if the patient has relapsed/refractory or metastatic disease AND the patient has tried one systemic chemotherapy regimen AND pt has Ewing sarcoma or osteosarcoma. Colon and Rectal cancer/Appendiceal cancer-approve if the patient has advanced or metastatic disease, has been previously treated with a fluoropyrimidine, oxaliplatin, irinotecan and if the patient meets one of the following (i or ii): i. patient's tumor or metastases are wild-type RAS (KRAS wild type and NRAS wild type), the patient has tried Erbitux or Vectibix or the patient's tumor did not originate on the left side of the colon (from the splenic fixture to rectum) or ii. the patient's tumor or metastases has a RAS mutation (either KRAS mutation or NRAS mutation). CNS tumors (Glioblastoma or H3-mutated high-grade glioma)-approve if the patient has recurrent or progressive disease. Uterine sarcoma- (A and B): A) pt has recurrent, advanced, inoperable, or metastatic disease, and B) tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Soft tissue Sarcoma, Bone Cancer, CNS tumors (Glioblastoma/H3-mutated high-grade glioma), Appendiceal cancer, Uterine sarcoma
Part B Prerequisite	No

SUCRAID

Products Affected

- SUCRAID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, genetic and lab test results (as specified in the Other Criteria field)
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a geneticist, gastroenterologist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of congenital diarrheal disorders
Coverage Duration	1 year
Other Criteria	Approve if the patient meets the following criteria (A and B): A) The diagnosis is established by one of the following (i or ii): i. Patient has endoscopic biopsy of the small bowel with disaccharidase levels consistent with congenital sucrose-isomaltase deficiency as evidenced by ALL of the following (a, b, c, and d): a) Decreased (usually absent) sucrase (normal reference: greater than 25 U/g protein), b) Decreased or normal isomaltase (palatinase) [normal reference: greater than 5 U/g protein], c) Decreased maltase (normal reference: greater than 100 U/g protein), d) Decreased or normal lactase (normal reference: greater than 15 U/g protein) OR ii. Patient has a molecular genetic test demonstrating homozygous or compound heterozygous pathogenic or likely pathogenic sucrase-isomaltase gene variant AND B) Patient has symptomatic congenital sucrose-isomaltase deficiency (e.g., diarrhea, bloating, abdominal cramping).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SUNITINIB

Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Gastrointestinal stromal tumors (GIST), approve if the patient has tried imatinib or Ayvakit or if the patient has succinate dehydrogenase (SDH)-deficient GIST. Chordoma, approve if the patient has recurrent disease. Differentiated thyroid carcinoma, approve if the patient meets (A or B): A) has papillary or follicular thyroid carcinoma and the disease is refractory to radioactive iodine therapy or B) has oncocytic (formerly Hurthle cell) carcinoma. Medullary thyroid carcinoma, approve if the patient has tried at least one systemic therapy. Meningioma, approve if the patient has recurrent or progressive disease. Thymic carcinoma - has tried at least one systemic chemotherapy. Renal Cell Carcinoma (RCC)- approve if the patient has relapsed or advanced disease. Neuroendocrine tumors of the pancreas-approve for advanced or metastatic disease. Pheochromocytoma/paraganglioma-approve if the patient has unresectable or metastatic disease. Myeloid/lymphoid neoplasms with eosinophilia-approve if the tumor has an FLT3 rearrangement.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chordoma, angiosarcoma, solitary fibrous tumor/hemangiopericytoma, alveolar soft part sarcoma (ASPS), extraskeletal myxoid chondrosarcoma, differentiated (ie, papillary, follicular, and oncocytic carcinoma) thyroid carcinoma, medullary thyroid carcinoma, meningioma, thymic carcinoma, pheochromocytoma/paraganglioma, myeloid/lymphoid neoplasms with

PA Criteria	Criteria Details
	eosinophilia, GIST-unresectable succinate dehydrogenase (SDH)-deficient GIST, or use after avapritinib.
Part B Prerequisite	No

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	Patients with unknown CFTR gene mutations, Combination therapy with Orkambi, Kalydeco or Trikafta
Required Medical Information	Diagnosis, specific CFTR gene mutations
Age Restrictions	Six years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	1 year
Other Criteria	CF - Approve if the pt meets A, B and C: A) pt has at least one mutation in the CFTR gene that is considered to be pathogenic or likely pathogenic or patient has TWO copies of the F508 del mutation, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SYMLIN

Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TABRECTA

Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-Approve if the patient has advanced or metastatic disease AND the tumor is positive for a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping or high-level MET amplification, as detected by an approved test.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-small cell lung cancer with high-level MET amplification.
Part B Prerequisite	No

Tadalafil

Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Indication for which tadalafil is being prescribed.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 mos.
Other Criteria	Benign prostatic hyperplasia (BPH), after confirmation that tadalafil is being prescribed as once daily dosing, to treat the signs and symptoms of BPH and not for the treatment of erectile dysfunction (ED).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TAFAMIDIS

Products Affected

- VYNDAMAX
- VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with Onpattro or Tegsedi or Wainua. Concurrent use of Vyndaqel and Vyndamax.
Required Medical Information	Diagnosis, genetic tests and lab results (as specified in the Other Criteria field)
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a physician who specializes in the treatment of amyloidosis
Coverage Duration	1 year
Other Criteria	Cardiomyopathy of Wild-Type or Hereditary Transthyretin Amyloidosis-approve if the diagnosis was confirmed by one of the following (i, ii or iii): i. A technetium pyrophosphate scan (i.e., nuclear scintigraphy), ii. Amyloid deposits are identified on cardiac biopsy OR iii. patient had genetic testing which, according to the prescriber, identified a TTR mutation AND Diagnostic cardiac imaging (e.g., echocardiogram, cardiac magnetic imaging) has demonstrated cardiac involvement (e.g., increased thickness of the ventricular wall or interventricular septum).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TAFINLAR

Products Affected

- TAFINLAR ORAL CAPSULE
- TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Tafinlar is being used. BRAF V600 mutations
Age Restrictions	1 year and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Melanoma with BRAF V600 mutation AND patient has unresectable, advanced (including Stage III or Stage IV disease) or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC, must have BRAF V600E mutation. Thyroid Cancer, anaplastic-must have BRAF V600-positive disease AND Tafinlar will be taken in combination with Mekinist, unless intolerant AND the patient has locally advanced or metastatic anaplastic disease. Thyroid Cancer, differentiated (i.e., papillary, follicular, or Hurthle cell) AND the patient has disease that is refractory to radioactive iodine therapy AND the patient has BRAF-positive disease. Biliary Tract Cancer-approve if the patient has tried at least one systemic chemotherapy regimen, patient has BRAF V600 mutation positive disease and the medication will be taken in combination with Mekinist. Central Nervous System Cancer-approve if the medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, or c): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma OR ii. Recurrent or progressive disease for one of the following conditions (a, b, c, or d): a) glioma OR b) Isocitrate dehydrogenase-2 (IDH2)-mutant astrocytoma OR c) Glioblastoma OR d) Oligodendrogloma OR iii. Melanoma with brain metastases AND patient has BRAF V600 mutation-positive disease AND

PA Criteria	Criteria Details
	<p>medication will be taken in combination with Mekinist (trametinib tablets). Histiocytic neoplasm-approve if patient has Langerhans cell histiocytosis and one of the following (a, b, or c): a) Multisystem disease OR b) Pulmonary disease OR c) Central nervous system lesions OR patient has Erdheim Chester disease AND patient has BRAF V600-mutation positive disease. Metastatic or solid tumors-approve if BRAF V600 mutation-positive disease AND medication will be taken in combination with Mekinist (trametinib tablets) AND patient has no satisfactory alternative treatment options. Ovarian, Fallopian Tube, or Primary Peritoneal Cancer-approve if the patient meets the following (A, B, and C): A) Patient has recurrent disease, AND B) Patient has BRAF V600 mutation-positive disease, AND C) The medication will be taken in combination with Mekinist (trametinib tablets). Hairy Cell Leukemia, approve if pt has not previously been treated with a BRAF inhibitor therapy and this will be used for relapsed/refractory disease and will be taken in combination with Mekinist. Small bowel adenocarcinoma, approve if pt has BRAF V600E mutation-positive advanced or metastatic disease and this will be used with Mekinist AND (i or ii): i) this will be used as initial therapy and pt has received previous FOLFOX/CAPEOX therapy in the adjuvant setting within the past 12 months or has a contraindication, or (ii) this will be used as second-line and subsequent therapy.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Differentiated Thyroid Cancer, Biliary tract cancer, central nervous system cancer, histiocytic neoplasm, Ovarian, Fallopian Tube, or Primary Peritoneal Cancer, hairy cell leukemia, small bowel adenocarcinoma
Part B Prerequisite	No

TAGRISSO

Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NSCLC-EGFR Mutation Positive (other than EGFR T790M-Positive Mutation)- approve if the patient has advanced or metastatic disease, has EGFR mutation-positive NSCLC as detected by an approved test. Note-examples of EGFR mutation-positive NSCLC include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S7681. NSCLC-EGFR T790M mutation positive- approve if the patient has advanced or metastatic EGFR T790M mutation-positive NSCLC as detected by an approved test and has progressed on treatment with at least one of the EGFR tyrosine kinase inhibitors. NSCLC- Post resection-approve if the patient has completely resected disease and has received previous adjuvant chemotherapy or if the patient is ineligible to receive platinum based chemotherapy and the patient has EGFR exon 19 deletions or exon 21 L858R substitution mutations, as detected by an approved test. NSCLC- Unresectable Stage III - approve if the patient has locally advanced, unresectable (stage III) disease AND EGFR exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an approved test AND not had disease progression during or following platinum-based chemoradiation therapy. (Note: Patients could have received concurrent or sequential chemoradiation therapy.)
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

TALZENNA

Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Recurrent or metastatic breast cancer-approve if the patient has germline BRCA mutation-positive disease. Prostate cancer - approve if the patient has metastatic castration resistant prostate cancer, AND is using this medication concurrently with a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy AND the patient has homologous recombination repair (HRR) gene-mutated disease [Note: HRR gene mutations include ATM, ATR, BRCA1, BRCA2, CDK12, CHEK2, FANCA, MLH1, MRE11A, NBN, PALB2, or RAD51C] AND the medication is used in combination with Xtandi (enzalutamide capsules and tablets).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TASIGNA

Products Affected

- *nilotinib hcl oral capsule 150 mg, 200 mg, 50 mg*
- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Tasigna is being used. For indication of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For indication of gastrointestinal stromal tumor (GIST) and melanoma, cutaneous, prior therapies tried. For melanoma, cutaneous, KIT mutation status.
Age Restrictions	GIST/Myeloid/lymphoid neoplasms/melanoma, cutaneous-18 years and older, ALL - 15 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Acute lymphoblastic leukemia, philadelphia chromosome positive-approve. CML, philadelphia chromosome positive or BCR::ABL1-mutation positive chronic myeloid leukemia- approve. For GIST, approve if the patient has tried two of the following: imatinib, avapritinib, sunitinib, dasatinib, regorafenib or ripretinib. Myeloid/lymphoid neoplasms with eosinophilia- approve if the tumor has an ABL1 rearrangement. Pigmented villonodular synovitis/tenosynovial giant cell tumor-approve if the patient has tried Turalio or Romvimza or cannot take Turalio or Romvimza, according to the prescriber. For melanoma, cutaneous - approve if the patient has metastatic or unresectable disease AND has an activating KIT mutation AND has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Philadelphia positive Acute Lymphoblastic Leukemia (ALL) and Gastrointestinal Stromal Tumor (GIST), Pigmented villonodular synovitis/tenosynovial giant cell tumor, Myeloid/Lymphoid neoplasms with Eosinophilia, melanoma cutaneous.

PA Criteria	Criteria Details
Part B Prerequisite	No

TAZAROTENE

Products Affected

- *tazarotene topical cream*
- *tazarotene topical gel*

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic uses
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TAZVERIK

Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Epithelioid Sarcoma-16 years and older, Follicular Lymphoma-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Epithelioid Sarcoma-approve if the patient has metastatic or locally advanced disease and the patient is not eligible for complete resection. Follicular Lymphoma-approve if the patient has relapsed or refractory disease and there are no appropriate alternative therapies or the patient has tried at least two prior systemic therapies.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TEPMETKO

Products Affected

- TPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NSCLC-approve if the patient has advanced or metastatic disease and the tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutations or patient has high-level MET amplification, as detected by an approved test.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-small cell lung cancer with high-level MET amplification.
Part B Prerequisite	No

TERIPARATIDE

Products Affected

- BONSITY
- *teriparatide (only ndcs starting with 47781)*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other medications for osteoporosis
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	High risk for fracture-2 yrs, Not high risk-approve a max of 2 yrs of therapy (total)/lifetime.
Other Criteria	<p>INITIAL THERAPY: Postmenopausal Osteoporosis (PMO) Treatment, Increase Bone Mass in Men (see Note 1 below) with Primary or Hypogonadal Osteoporosis, and Treatment of Glucocorticosteroid-Induced Osteoporosis (GIO): (one of A, B, C, D or E): A) tried one oral bisphosphonate or cannot take due to swallowing difficulties or inability to remain upright after administration, B) pre-existing gastrointestinal condition (e.g., esophageal lesions/ulcers, abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), C) tried an IV bisphosphonate (PMO-ibandronate or zoledronic acid, all other diagnoses-zoledronic acid), D) severe renal impairment (creatinine clearance [CrCL] less than 35 mL/min) or chronic kidney disease (CKD), or E) patient had an osteoporotic fracture or fragility fracture at any time in the past.</p> <p>CONTINUATION THERAPY: ALL INDICATIONS: if the patient has taken teriparatide for two years, approve if the patient is at high risk for fracture. Examples of high risk for fracture include a previous osteoporotic fracture or fragility fracture, receipt of medications that increase the risk of osteoporosis, advanced age, and very low bone mineral density. Note 1: a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.</p>
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

TETRABENAZINE

Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	For treatment of chorea associated with Huntington's disease, Tourette syndrome or related tic disorders, hyperkinetic dystonia, or hemiballism, must be prescribed by or after consultation with a neurologist. For TD, must be prescribed by or after consultation with a neurologist or psychiatrist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Chorea associated with Huntington's Disease-approve if the diagnosis of Huntington's Disease is confirmed by genetic testing.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Tardive dyskinesia (TD). Tourette syndrome and related tic disorders. Hyperkinetic dystonia. Hemiballism.
Part B Prerequisite	No

THALOMID

Products Affected

- THALOMID ORAL CAPSULE 100 MG,
50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	MM, myelofibrosis, histiocytic neoplasms-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Erythem Nodosum Leprosum-approve. Multiple Myeloma-approve. Discoid lupus erythematosus or cutaneous lupus erythematosus, approve if the patient has tried at least two other therapies (eg, corticosteroids [oral, topical, intralesional], hydroxychloroquine, tacrolimus [Protopic], methotrexate, dapsone, acitretin [Soriatane]). Myelofibrosis, approve if according to the prescriber the patient has anemia and has serum erythropoietin levels greater than or equal to 500 mU/mL or if the patient has serum erythropoietin level less than 500 mU/mL and experienced no response or loss of response to erythropoietic stimulating agents. Prurigo nodularis, approve. Recurrent aphthous ulcers or aphthous stomatitis, approve if the patient has tried at least two other therapies (eg, topical or intralesional corticosteroids, systemic corticosteroids, topical anesthetics/analgesics [eg, benzocaine lozenges], antimicrobial mouthwashes [eg, tetracycline], acyclovir, colchicine). Kaposi's Sarcoma-approve if the patient has tried at least one regimen or therapy and has relapsed or refractory disease. Castleman's disease-approve if the patient has multicentric Castleman's disease and is negative for the human immunodeficiency virus (HIV) and human herpesvirus-8 (HHV-8). Histiocytic neoplasms-approve if pt has Langerhans cell histiocytosis with single-system multifocal skin disease or Rosai-Dorfman cutaneous disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Discoid lupus erythematosus or cutaneous lupus erythematosus, Myelofibrosis, Prurigo nodularis, Recurrent aphthous ulcers or aphthous stomatitis, Kaposi's Sarcoma, Castleman's Disease, histiocytic neoplasms.
Part B Prerequisite	No

TIBSOVO

Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, IDH1 Status
Age Restrictions	All diagnoses (except chondrosarcoma)-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML- approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive, as detected by an approved test. Cholangiocarcinoma-approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive and has been previously treated with at least one chemotherapy regimen (Part B before Part D Step Therapy - applies only to beneficiaries enrolled in an MA-PD plan). Chondrosarcoma-approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive. Central nervous system cancer-approve if the patient has oligodendrogloma or astrocytoma. Myelodysplastic Syndrome-approve if patient has isocitrate dehydrogenase-1 (IDH1) mutation-positive disease AND relapsed or refractory disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chondrosarcoma, Central nervous system cancer
Part B Prerequisite	Yes

TOBRAMYCIN (NEBULIZATION)

Products Affected

- *tobramycin in 0.225 % nacl*
- *tobramycin inhalation*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Bronchiectasis, Non-cystic fibrosis-18 years and older
Prescriber Restrictions	CF-prescr/consult w/pulm/phys specializes in tx of CF.Bronchiectasis, non CF-prescr/consult w/pulm
Coverage Duration	1 year
Other Criteria	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Cystic fibrosis/Bronchiectasis, non-cystic fibrosis-approve if the patient has pseudomonas aeruginosa in the culture of the airway.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Bronchiectasis, non-cystic fibrosis
Part B Prerequisite	No

TOLVAPTAN

Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Jynarque.
Required Medical Information	Serum sodium less than 125 mEq/L at baseline or less marked hyponatremia, defined as serum sodium less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion).
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 30 days for initial therapy, 3 months for continuation of therapy
Other Criteria	Hyponatremia, initial therapy (including new starts, patients on therapy for less than 30 days, and patients restarting therapy) - Pt must meet ONE of the following: 1. serum sodium less than 125 mEq/L at baseline, OR 2. marked hyponatremia, defined as less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion), OR 3. patient has already been started on tolvaptan and has received less than 30 days of therapy. Hyponatremia, continuation of therapy for patients established on therapy for at least 30 days - approve if the serum sodium level has increased from baseline (prior to initiating the requested drug) OR if the patient experienced improvement in at least one symptom, such as nausea, vomiting, headache, lethargy, or confusion.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOPICAL AGENTS FOR ATOPIC DERMATITIS

Products Affected

- *pimecrolimus*
- *tacrolimus topical*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Authorize use in patients who have tried a prescription strength topical corticosteroid (brand or generic) for the current condition. Dermatologic condition on or around the eyes, eyelids, axilla, or genitalia, authorize use without a trial of a prescription strength topical corticosteroid.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOPICAL RETINOID PRODUCTS

Products Affected

- *tretinoin topical*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for cosmetic use.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOPIRAMATE/ZONISAMIDE

Products Affected

- EPRONTIA
- *topiramate oral capsule, sprinkle 15 mg, 25 mg*
- *topiramate oral solution*
- *topiramate oral tablet*
- ZONISADE
- *zonisamide*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for weight loss or smoking cessation.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TRANSDERMAL FENTANYL

Products Affected

- *fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr*

PA Criteria	Criteria Details
Exclusion Criteria	Acute (i.e., non-chronic) pain.
Required Medical Information	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been tried and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, sickle cell disease, in hospice or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids (including transdermal fentanyl products) require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

TRELSTAR

Products Affected

- TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prostate cancer: Prescribed by or in consultation with a oncologist or urologist. Head and neck cancer - salivary gland tumors: Prescribed by or in consultation with a oncologist.
Coverage Duration	1 year
Other Criteria	Part B versus Part D determination will be made at time of prior authorization review per CMS guidance. Prostate cancer: Patients new to therapy, are required to try Eligard or Orgovyx prior to approval of Trelstar. Head and neck cancer - salivary gland tumors: approve if patient has recurrent, unresectable, or metastatic disease and androgen receptor-positive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Head and neck cancer - salivary gland tumors
Part B Prerequisite	No

TREMFYA

Products Affected

- TREMFYA ONE-PRESS
- TREMFYA PEN INDUCTION PK-CROHN
- TREMFYA PEN SUBCUTANEOUS PEN INJECTOR 200 MG/2 ML
- TREMFYA SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
Required Medical Information	Diagnosis
Age Restrictions	PP/UC/CD- 18 years of age and older (initial therapy)
Prescriber Restrictions	PP-Prescribed by or in consultation with a dermatologist (initial therapy only). PsA-prescribed by or in consultation with a dermatologist or rheumatologist (initial therapy). UC/CD-prescribed by or in consultation with a gastroenterologist (initial therapy).
Coverage Duration	Approve through end of plan year
Other Criteria	PP, intial therapy - approve if the pt meets one of the following criteria: 1) pt has tried at least one traditional systemic agent (eg, MTX, cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (note: a biologic that is not a biosimilar of the requested product will also count) OR 2) pt has a contraindication to MTX as determined by the prescribing physician. ULCERATIVE COLITIS- pt will receive 3 induction doses with Tremfya IV within 3 months of initiating Tremfya SC. CROHN'S DISEASE (CD) [one of A, B, C, or D]: A) tried or currently taking corticosteroid (CS) or CS is contraindicated, B) tried one other conventional systemic therapy (e.g., azathioprine, 6-mercaptopurine [6-MP], methotrexate [MTX], certolizumab, infliximab, ustekinumab, vedolizumab), C) had ileocolonic resection, or D) enterocutaneous (perianal or abdominal) or rectovaginal fistulas. PP/PsA/UC/CD continuation of therapy - approve if the pt is responding to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

TRIENTINE

Products Affected

- *trientine oral capsule 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician.
Coverage Duration	Authorization will be for 1 year
Other Criteria	For Wilson's Disease, approve if the patient meets A and B: A) Diagnosis of Wilson's disease is confirmed by ONE of the following (i or ii): i. Genetic testing results confirming biallelic pathogenic ATP7B mutations (in either symptomatic or asymptomatic individuals), OR ii. Confirmation of at least two of the following (a, b, c, or d): a. Presence of Kayser-Fleischer rings, OR b. Serum ceruloplasmin levels less than 20mg/dL, OR c. Liver biopsy findings consistent with Wilson's disease, OR d. 24-hour urinary copper greater than 40 micrograms/24 hours, AND B) Patient meets ONE of the following: 1) Patient has tried a penicillamine product and per the prescribing physician the patient is intolerant to penicillamine therapy, OR 2) Per the prescribing physician, the patient has clinical features indicating the potential for intolerance to penicillamine therapy (ie, history of any renal disease, congestive splenomegaly causing severe thrombocytopenia, autoimmune tendency), OR 3) Per the prescribing physician, the patient has a contraindication to penicillamine therapy, OR 4) The patient has neurologic manifestations of Wilson's disease, OR 5) The patient is pregnant, OR 6) the patient has been started on therapy with trientine.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

TRIKAFTA

Products Affected

- TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL
- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	Patients with unknown CFTR gene mutations. Combination therapy with Orkambi, Kalydeco or Symdeko.
Required Medical Information	Diagnosis, specific CFTR gene mutations, concurrent medications
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	1 year
Other Criteria	CF - Approve if the pt meets A, B and C: A) pt has at least one mutation in the CTFR gene that is considered to be pathogenic or likely pathogenic, and B) pt must have positive CF newborn screening test or family history of CF or clinical presentation consistent with signs and symptoms of CF and C) evidence of abnormal CFTR function as demonstrated by i, ii, or iii: (i) elevated sweat chloride test or (ii) two CFTR mutations or (iii) abnormal nasal potential difference.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TRUQAP

Products Affected

- TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast Cancer-Approve if the patient meets the following (A, B, C, D and E): A) Patient has locally advanced or metastatic disease, AND B) Patient has hormone receptor positive (HR+) disease, AND C) Patient has human epidermal growth factor receptor 2 (HER2)-negative disease, AND D) Patient has at least one phosphatidylinositol 3-kinase (PIK3CA), serine/threonine protein kinase (AKT1), or phosphatase and tensin homolog (PTEN)-alteration, AND E) Patient meets one of the following (i or ii): i. Patient has had progression with at least one endocrine-based regimen in the metastatic setting (Note: Examples of endocrine therapy include anastrozole, exemestane, and letrozole.) and has had progression with at least one cyclin-dependent kinase (CDK) 4/6 inhibitor in the metastatic setting (Note: Examples of CDK4/6 inhibitor include: Ibrance (palbociclib tablets or capsules), Verzenio (abemaciclib tablets), Kisqali (ribociclib tablets), Kisqali Femara Co-Pack (ribociclib and letrozole tablets) OR ii. Patient has recurrence on or within 12 months of completing adjuvant endocrine therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TUKYSA

Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast Cancer-approve if the patient has recurrent or metastatic human epidermal growth factor receptor 2 (HER2)-positive disease, has received at least one prior anti-HER2-based regimen in the metastatic setting and Tukysa is used in combination with trastuzumab and capecitabine. Colon/Rectal Cancer-approve if the requested medication is used in combination with trastuzumab, patient has unresectable or metastatic disease, human epidermal growth factor receptor 2 (HER2)-amplified disease, AND Patient's tumor or metastases are wild-type RAS (KRAS wild-type and NRAS wild-type). Biliary tract cancer- approve if the patient meets all of (a, b, c, and d): a) unresectable or metastatic disease, b) HER2 positive disease, c) tried at least one systemic regimen, d) will use in combination with trastuzumab.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Biliary tract cancer
Part B Prerequisite	No

TURALIO

Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Tenosynovial Giant Cell Tumor (Pigmented Villonodular Synovitis)- approve if, according to the prescriber, the tumor is not amenable to improvement with surgery. Histiocytic Neoplasms-approve if the patient has a colony stimulating factor 1 receptor (CSF1R) mutation AND has one of the following conditions (i, ii, or iii): i. Langerhans cell histiocytosis OR ii. Erdheim-Chester disease OR iii. Rosai-Dorfman disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Histiocytic Neoplasms
Part B Prerequisite	No

UBRELVY

Products Affected

- UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Migraine, Acute treatment-approve
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

UPTRAVI

Products Affected

- UPTRAVI ORAL TABLET
- UPTRAVI ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with Other Inhaled or Parenteral Prostacyclin Agents Used for Pulmonary Hypertension.
Required Medical Information	Confirmation of right heart catheterization, medication history (as described in Other Criteria)
Age Restrictions	N/A
Prescriber Restrictions	PAH must be prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	1 year
Other Criteria	Must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1). Patient new to therapy must meet a) OR b): a) tried one or is currently taking one oral therapy for PAH for 30 days, unless patient has experienced treatment failure, intolerance, or oral therapy is contraindicated: PDE5 inhibitor (eg, sildenafil, Revatio), endothelin receptor antagonist (ERA) [eg, Tracleer, Letairis or Opsumit], or Adempas, OR b) receiving or has received in the past one prostacyclin therapy for PAH (eg, Orenitram, Ventavis, or epoprostenol injection).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VALCHLOR

Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Cutaneous lymphoma-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Cutaneous Lymphomas (Note-includes mycosis fungoides/Sezary syndrome, primary cutaneous B-cell lymphoma, primary cutaneous CD30+ T-cell lymphoproliferative disorders)-approve. Adult T-Cell Leukemia/Lymphoma-approve if the patient has chronic/smoldering subtype of adult T-cell leukemia/lymphoma. Langerhans cell histiocytosis-approve if the patient has unifocal Langerhans cell histiocytosis with isolated skin disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Adults with T-cell leukemia/lymphoma, Langerhans Cell Histiocytosis
Part B Prerequisite	No

VALTOCO

Products Affected

- VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications used at the same time
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiseizure medication(s).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VANCOMYCIN

Products Affected

- *vancomycin oral capsule 125 mg, 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	2 weeks
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VANFLYTA

Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Acute Myeloid Leukemia: approve if the patient has FLT3-ITD mutation-positive disease as detected by an approved test and this medication is being used for induction, re-induction, consolidation, or maintenance treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VENCLEXTA

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy
Age Restrictions	18 years and older (all diagnoses except ALL)
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML-approve if used in combination with azacitidine, decitabine, or cytarabine. CLL/SLL- approve. ALL- approve if relapsed/refractory disease and will be used in combination with chemotherapy. Hairy cell leukemia- approve if disease resistance to BRAF inhibitor therapy. Mantle Cell Lymphoma- approve if (A or B): A) the patient has tried at least one systemic regimen or B) patient has TP53 mutation and will use this as induction therapy in combination with Brukinsa (zanubrutinib) and Gazyva (obinutuzumab intravenous infusion). MDS- approve if pt meets (A and B): A) pt meets (i or ii): (i) has chronic myelomonocytic leukemia-2 or (ii) has higher risk disease (note: includes international prognostic scoring system (IPSS-R) intermediate-, high-, or very-high risk disease) and B) will use in combination with azacitidine or decitabine. Myeloproliferative neoplasm- approve if pt has accelerated or blast phase disease and will use in combination with azacitidine or decitabine. Multiple Myeloma- approve if the patient has t (11,14) translocation AND has tried at least one systemic regimen for multiple myeloma AND Venclexta will be used in combination with dexamethasone. Systemic light chain amyloidosis-approve if the patient has t (11, 14) translocation and has tried at least one systemic regimen. Waldenstrom Macroglobulinemia/Lymphoplasmacytic Lymphoma-approve if the patient has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Mantle Cell Lymphoma, waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma, multiple myeloma, systemic light chain amyloidosis, acute lymphoblastic leukemia, hairy cell leukemia, myelodysplastic syndrome, myeloproliferative neoplasm
Part B Prerequisite	No

VERZENIO

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: HR status, HER2 status, previous medications/therapies tried, concomitant therapy, menopausal status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Breast Cancer, Early-pt meets (A,B,C and D): A)Pt HR+disease, AND B) HER2-negative breast cancer, AND C)node-positive disease at high risk of recurrence AND D)meets 1 of the following (i or ii): i.Verzenio will be used in combo w/anastrozole, exemestane, or letrozole AND pt meets one of the following (a,b, or c): a)Pt is postmenopausal woman, OR b)Pt is pre/perimenopausal woman and meets one of the following 1 or 2:1-Pt receiving ovarian suppression/ablation with a GnRH agonist, OR 2-Pt had surgical bilateral oophorectomy or ovarian irradiation, OR c)Pt is man and pt is receiving a GnRH analog, OR ii.Verzenio will be used in combo with tamoxifen AND pt meets 1 of the following (a or b): a)Pt is postmenopausal woman or man OR b)Pt is pre/perimenopausal woman and meets 1 of the following 1 or 2:1-Pt is receiving ovarian suppression/ablation with a GnRH agonist, OR 2-Pt had surgical bilateral oophorectomy or ovarian irradiation. Breast Cancer-Recurrent or Metastatic in Women-pt meets (A, B and C): A)has HR+ disease, AND B)Pt meets 1 of following criteria (i or ii): i.Pt is postmenopausal woman, OR ii.Pt is pre/perimenopausal woman and meets 1 of the following (a or b): a)receiving ovarian suppression/ablation with a GnRH agonist, OR b)Pt had surgical bilateral oophorectomy or ovarian irradiation, AND C) either (1 or 2): 1) HER2-negative breast cancer and Pt meets 1 of following criteria (i, ii, or iii): i.Verzenio will be used in combo with anastrozole, exemestane, or letrozole, OR ii.Verzenio will be used in combo with

PA Criteria	Criteria Details
	<p>fulvestrant, OR iii.pt meets following conditions (a, b, and c): a)Verzenio will be used as monotherapy, AND b)Pt's breast cancer has progressed on at least 1 prior endocrine therapy, AND c)has tried chemo for metastatic breast cancer or 2)has HER2-positive breast cancer and has received at least 3 prior anti-HER2-based regimens in metastatic setting and will use this in combo with fulvestrant and trastuzumab.Breast Cancer-Recurrent or Metastatic in Men-pt meets following criteria (A and B): A)HR+ disease, AND B)either (1 or 2): 1) HER2-negative disease and Pt meets 1 of following criteria (i, ii, or iii): i.Pt meets BOTH of the following conditions (a and b): a)receiving a GnRH analog, AND b)Verzenio will be used in combo with anastrozole, exemestane, or letrozole, OR ii.Verzenio will be used in combo with fulvestrant, OR iii.Pt meets the following conditions (a, b, and c): a)Verzenio will be used as monotherapy, AND b)Pt's breast cancer has progressed on at least 1 prior endocrine therapy, AND c)Pt has tried chemo for metastatic breast cancer, or 2) has HER2-positive disease and has received at least 3 prior anti-HER2-based regimen in metastatic setting and will use this medication in combo with fulvestrant and trastuzumab. Endometrial cancer-pt meets all of (A, B, And C): A)has recurrent or metastatic disease, and B)has estrogen receptor (ER)-positive tumors, and C)will be using in combination with letrozole.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Endometrial cancer
Part B Prerequisite	No

VIGABATRIN

Products Affected

- *vigabatrin*
- *vigadron*
- *vigpoder*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medication history (complex partial seizures)
Age Restrictions	Refractory complex partial seizures - patients 2 years of age or older. Infantile spasms - patients less than or equal to 2 years of age
Prescriber Restrictions	Prescribed by, or in consultation with, a neurologist
Coverage Duration	Infantile spasms- 6 months. Treatment-Refractory Partial Seizures- initial 3 months, cont 1 year
Other Criteria	Infantile spasms-requested medication is being used as monotherapy. Treatment refractory complex partial seizures intial-patient has tried and/or is concomitantly receiving at least two other antiepileptic drugs. Treatment refractory complex partial seizures continuation- the patient is responding to therapy (e.g., reduced seizure severity, frequency, and/or duration).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VITRAKVI

Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, NTRK gene fusion status
Age Restrictions	Pediatric Diffuse High-Grade Glioma- less than or equal to 21 years old
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Solid tumors - approve if the tumor is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND the tumor is metastatic or surgical resection of tumor will likely result in severe morbidity. Pediatric diffuse high grade glioma - approve if (A and B): A) tumor is positive for NTRK gene fusion and B) meets (i or ii): i) medication is used as adjuvant therapy or ii) medication is used for recurrent or progressive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pediatric Diffuse High-Grade Glioma
Part B Prerequisite	No

VIZIMPRO

Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, EGFR status, exon deletions or substitutions
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	NSCLC-approve if the patient has advanced or metastatic disease, has sensitizing EGFR mutation-positive NSCLC as detected by an approved test. Note: Examples of sensitizing EGFR mutation-positive NSCLC include the following mutations: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X and S7681.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VONJO

Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Myelofibrosis (MF), including primary MF, post-polycythemia Vera MF, and Post-Essential Thrombocythemia MF-approve if the patient meets either (A, B, or C): (A) the patient has a platelet count of less than $50 \times 10^9 /L$ (less than 50,000/mcL) OR (B) Patient has a platelet count of greater than or equal to $50 \times 10^9 /L$ (greater than or equal to 50,000/mcL) and has high-risk disease, OR (C) patient has myelofibrosis-associated anemia. Accelerated or blast phase myeloproliferative neoplasm- approve if the patient has at least one disease- related symptom (Examples: fatigue, fever, night sweats, weight loss, abdominal discomfort, splenomegaly, thrombocytosis, or leukocytosis).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Accelerated or blast phase myeloproliferative neoplasm
Part B Prerequisite	No

VORANIGO

Products Affected

- VORANIGO ORAL TABLET 10 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	12 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	GLIOMAS-All of (A, B and C): A. Susceptible isocitrate dehydrogenase-1 (IDH1) or IDH2 mutation-positive disease, AND B. Grade 2 oligodendrolioma OR Grade 2 astrocytoma, AND C. Prior surgery, including biopsy, sub-total resection, or gross total resection
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VORICONAZOLE (ORAL)

Products Affected

- *voriconazole*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Aspergillus-Prophy, systemic w/risk neutropenia-Prophy, systemic w/HIV-Prophy/Tx-6 mo, others-3 mo
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Aspergillus Infections - prophylaxis, oropharyngeal candidiasis (fluconazole-refractory) - treatment, candida endophthalmitis - treatment, blastomycosis - treatment, fungal infections (systemic) in patients at risk of neutropenia - prophylaxis, fungal infections (systemic) in patients with human immunodeficiency virus (HIV) - prophylaxis or treatment.
Part B Prerequisite	No

VOSEVI

Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
Coverage Duration	Will be c/w AASLD guidance and inclusive of treatment already received for the requested drug
Other Criteria	Criteria will be applied consistent with current AASLD/IDSA guidance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance
Part B Prerequisite	No

VOTRIENT

Products Affected

- *pazopanib*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Soft tissue sarcoma other than GIST-approve if the patient has advanced or metastatic disease and has ONE of the following: alveolar soft part sarcoma, angiosarcoma, desmoid tumors (aggressive fibromatosis, dermatofibrosarcoma protuberans with fibrosaromatous transformation, non-adipocytic sarcoma or pleomorphic rhabdomyosarcoma. Differentiated (ie, papillary, follicular, Hurthle) thyroid carcinoma, approve if the patient is refractory to radioactive iodine therapy. Uterine sarcoma, approve if the patient has recurrent or metastatic disease. Renal Cell Carcinoma, Clear Cell or non-Clear Cell histology-approved if the patient has relapsed or advanced disease or VonHippel-Lindau disease. Ovarian Cancer (ie, Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer) - approve if the patient has persistent or recurrent disease. GIST - approve if the patient has succinate dehydrogenase (SDH)-deficient GIST OR the patient has tried TWO of the following: Gleevec, Ayvakit, Sutent, Sprycel, Qinlock or Stivarga. Medullary Thyroid Carcinoma, approve if the patient has tried at least one systemic therapy. Bone cancer-approve if the patient has chondrosarcoma and has metastatic widespread disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (ie, papillary, follicular, oncocytic carcinoma) thyroid carcinoma. Uterine sarcoma, Epithelial Ovarian, Fallopian Tube, or

PA Criteria	Criteria Details
	Primary Peritoneal Cancer, Gastrointestinal Stromal Tumor (GIST), Medullary thyroid carcinoma, bone cancer.
Part B Prerequisite	No

VOWST

Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	30 days
Other Criteria	Prevention of recurrence of clostridioides difficile infection (CDI)-approve if the patient has completed a bowel prep (or will start a bowel prep the day before and at least 8 hours prior to taking the first dose), will not eat or drink for at least 8 hours prior to the first dose and will complete their antibacterial treatment for recurrent CDI 2-4 days before initiating treatment with Vowst and Vowst will not be used for the TREATMENT of CDI.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VUMERITY

Products Affected

- VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS)
Required Medical Information	Relapsing form of Multiple Sclerosis (MS), to include clinically-isolated syndrome, relapsing-remitting disease, and active secondary progressive disease
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of MS.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VYVGART

Products Affected

- VYVGART HYTRULO
SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant Use with Another Neonatal Fc Receptor Blocker, a Complement Inhibitor, or a Rituximab Product
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (CIDP: initial only, GMG: initial and continuation)
Prescriber Restrictions	Prescribed by or in consultation with a neurologist (CIDP: initial only, GMG: initial and continuation)
Coverage Duration	Initial-6 months, Continuation-1 year
Other Criteria	CIDP (Vyvgart Hytrulo only), Initial Therapy - approve if diagnosis was supported by electrodiagnostic studies AND pt previously received treatment with an intravenous or subcutaneous immune globulin and had inadequate efficacy or significant intolerance or patient has a contraindication to IV or SC immune globulin.. Generalized myasthenia gravis, Initial Therapy-Approve if the patient meets the following criteria (A, B, and C): A. Patient has confirmed anti-acetylcholine receptor antibody positive generalized myasthenia gravis, AND B. Patient has evidence of unresolved symptoms of generalized myasthenia gravis, such as difficulty swallowing, difficulty breathing, or a functional disability resulting in the discontinuation of physical activity (e.g., double vision, talking, impairment of mobility) AND C. patient has myasthenia gravis foundation of america classification of II to IV. CIDP (Vyvgart Hytrulo only), Cont therapy - pt has clinically significant improvement in neurologic symptoms (Examples include improvement in disability: nerve conduction study results improved or stabilized, physical examination shows improvement in neurological symptoms, strength, and sensation). Generalized myasthenia gravis, Continuation Therapy-Approve if patient is continuing to derive benefit from Vyvgart. For gMG: All treatment cycles should be no more frequent than every 50 days from the start of the previous treatment cycle.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

WELIREG

Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Pheochromocytoma/paraganglioma-12 years and older, Other indications-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Pheochromocytoma/paraganglioma- approve if pt has locally advanced, unresectable, or metastatic disease. Renal Cell Carcinoma- approve if pt has advanced disease AND has tried at least one programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor AND has tried at least one a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI). [Note: Examples of PD-1 inhibitor or PD-L1 inhibitor include: Keytruda (pembrolizumab intravenous infusion), Opdivo (nivolumab intravenous infusion), and Brevail (avelumab intravenous infusion). Examples of VEGF-TKI include Cabometyx (cabozantinib tablets), Lenvima (lenvatinib capsules), Inlyta (axitinib tablets), Fotivda (tivozanib capsules), pazopanib, sunitinib, and sorafenib.] Von Hippel-Lindau Disease-approve if the patient meets the following (A, B, and C): A) Patient has a von Hippel-Lindau (VHL) germline alteration as detected by genetic testing, B) Does not require immediate surgery and C) Patient requires therapy for ONE of the following conditions (i, ii, iii, or iv): i. Central nervous system hemangioblastomas, OR ii. Pancreatic neuroendocrine tumors, OR iii. Renal cell carcinoma, OR iv. Retinal hemangioblastoma.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

XALKORI

Products Affected

- XALKORI ORAL CAPSULE
- XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Anaplastic large cell lymphoma/IMT-patients greater than or equal to 1 year of age. All other diagnoses-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Metastatic non-small cell lung cancer-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease, as detected by an approved test and patients new to therapy must have a trial of Alecensa or Lorbrena prior to approval of Xalkori. Metastatic non-small cell lung cancer, approve if the patient has ROS1 rearrangement positive disease, as detected by an approved test. Anaplastic Large Cell Lymphoma-approve if the patient has anaplastic lymphoma kinase (ALK)-positive disease AND (i or ii): (i) the medication is used for palliative-intent therapy, or (ii) pt has relapsed or refractory disease. Histiocytic neoplasm-approve if the patient has ALK rearrangement/fusion-positive disease and meets one of the following criteria (i, ii, or iii): (i. Patient has Langerhans cell histiocytosis, OR ii. Patient has Erdheim-Chester disease OR iii. Patient has Rosai-Dorfman disease. NSCLC with MET mutation-approve if the patient has high level MET amplification or MET exon 14 skipping mutation. Inflammatory Myofibroblastic Tumor-approve if the patient has ALK positive disease and the patient has advanced, recurrent or metastatic disease or the tumor is inoperable. Melanoma, cutaneous-approve if the patient has ALK fusion disease or ROS1 fusion disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	NSCLC with high level MET amplification or MET Exon 14 skipping mutation, Histiocytic neoplasms, melanoma, cutaneous.
Part B Prerequisite	No

XDEMVY

Products Affected

- XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XELJANZ

Products Affected

- XELJANZ ORAL SOLUTION
- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a Targeted Synthetic DMARD for an inflammatory condition (eg, tocilizumab, anakinra, abatacept, rituximab, certolizumab pegol, etanercept, adalimumab, infliximab, golimumab). Concurrent use with potent immunosuppressants that are not methotrexate (MTX) [eg, azathioprine, tacrolimus, cyclosporine, mycophenolate mofetil].
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	AS/PsA/RA/UC-18 years and older (initial therapy)
Prescriber Restrictions	RA, JIA/JRA/AS prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or a dermatologist. UC-prescribed by or in consultation with a gastroenterologist. (all apply to initial therapy only)
Coverage Duration	Approve through end of plan year
Other Criteria	RA initial-approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. PsA initial, approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. UC-Approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least ONE tumor necrosis factor inhibitor for ulcerative colitis or was unable to tolerate a 3-month trial. Juvenile Idiopathic Arthritis (JIA) [or Juvenile Rheumatoid Arthritis] (regardless of type of onset) [Note: This includes patients with juvenile spondyloarthropathy/active sacroiliac arthritis]-initial-approve Xeljanz immediate release tablets or solution if the patient meets the following: patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. AS-approve Xeljanz/XR tablets if the patient has had a 3 month trial of at least one tumor necrosis factor inhibitor or was unable to tolerate a 3 month trial. Continuation Therapy - Patient must have responded, as determined by the prescriber.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XERMELO

Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapy, concomitant therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Initial therapy - approve if the patient meets ALL of the following criteria: 1) patient has been on long-acting somatostatin analog (SSA) therapy (eg, Somatuline Depot [lanreotide for injection]) AND 2) while on long-acting SSA therapy (prior to starting Xermelo), the patient continues to have at least four bowel movements per day, AND 3) Xermelo will be used concomitantly with a long-acting SSA therapy. Continuation therapy - approve if the patient is continuing to take Xermelo concomitantly with a long-acting SSA therapy for carcinoid syndrome diarrhea.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XIFAXAN

Products Affected

- XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Traveler's diarrhea - 12 years of age or older. Hepatic encephalopathy, irritable bowel syndrome with diarrhea - 18 years of age or older.
Prescriber Restrictions	Pouchitis - prescribed by or in consultation with a gastroenterologist
Coverage Duration	Enceph-6 mo, IBS w/diarrhea-14 days, TD-3 days, intest bact overgrowth-14 days, Pouchitis - 1 year
Other Criteria	Hepatic Encephalopathy-approve Xifaxan 550 mg tablets if the patient has previously had overt hepatic encephalopathy and the requested medication will be used concomitantly with lactulose, unless the patient has a contraindication or significant intolerance to treatment with lactulose. Irritable bowel syndrome with diarrhea-approve Xifaxan 550 mg tablets. Travelers Diarrhea-approve Xifaxan 200 mg tablets if the patient is afebrile and does not have blood in the stool. Small intestine bacterial overgrowth-approve Xifaxan 200mg or 550 mg tablets if the diagnosis has been confirmed by a glucose hydrogen breath test, lactulose hydrogen breath test, or small bowel aspiration and culture. Chronic antibiotic-dependent pouchitis- approve Xifaxan 200mg or 550mg tablets if patient meets all of (a, b, c and d): a) recurrent pouchitis (Note: recurrent pouchitis is typically considered history of at least 3 pouchitis episodes within a 12 month period), and b) episodes of pouchitis respond to antibiotic therapy but relapse shortly after antibiotic discontinuation, and c) alternative causes of recurrent pouchitis have been ruled out, and d) has tried long-term antibiotic therapy trials (at least 4 weeks) of BOTH ciprofloxacin and metronidazole for remission maintenance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Small intestine bacterial overgrowth, chronic antibiotic-dependent pouchitis
Part B Prerequisite	No

XOLAIR

Products Affected

- XOLAIR SUBCUTANEOUS AUTO-
INJECTOR 150 MG/ML, 300 MG/2 ML,
75 MG/0.5 ML
- XOLAIR SUBCUTANEOUS RECON
SOLN
- XOLAIR SUBCUTANEOUS SYRINGE
150 MG/ML, 300 MG/2 ML, 75 MG/0.5
ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another monoclonal antibody therapy.
Required Medical Information	Diagnosis
Age Restrictions	Moderate to severe persistent asthma-6 years and older. CIU-12 years and older. Polyps-18 years and older. Food Allergy-1 yr and older
Prescriber Restrictions	Moderate to severe persistent asthma if prescribed by, or in consultation with an allergist, immunologist, or pulmonologist. CIU if prescribed by or in consultation with an allergist, immunologist, or dermatologist. Polyps-prescribed by or in consult with an allergist, immunologist, or otolaryngologist. Food allergy- allergist or immunologist
Coverage Duration	asthma/CIU-Initial tx 4 months, Polyps-initial-6 months, continued tx 12 months, Food allergy-1 yr
Other Criteria	<p>MODERATE TO SEVERE PERSISTENT ASTHMA (A, B, C and D): A) baseline IgE greater than or equal to 30 IU/mL, and B) baseline positive skin test or in vitro test for 1 or more perennial or seasonal aeroallergens C) received at least 3 months of combination therapy with an inhaled corticosteroid (ICS) and additional asthma controller/maintenance medication (e.g., LABA, LAMA, leukotriene receptor antagonist, monoclonal antibody) [see Exception 1 below] and D) asthma is uncontrolled or was uncontrolled prior to receiving Xolair or another monoclonal antibody and meets one of the following (a, b, c, d, or e): a) experienced two or more asthma exacerbations requiring systemic CSs in the past year, b) experienced one or more asthma exacerbation requiring hospitalization/urgent care visit/emergency department visit in the past year, c) forced expiratory volume in 1 second (FEV1) less than 80% predicted, d) FEV1/forced vital capacity (FVC) less than 0.80, or e) asthma worsens upon tapering of oral CS.</p> <p>CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRwNP) [all of A, B, C, D, and E]: A) diagnosis by</p>

PA Criteria	Criteria Details
	<p>direct exam, endoscopy, or sinus CT scan, B) baseline (prior to Xolair or another monoclonal antibody that may lower IgE) IgE at least 30 IU/ml, C) at least two of the following symptoms for 6 months: nasal congestion, obstruction, discharge, reduction/loss of smell, D) tried intranasal CS and will continue in combination with Xolair, and E) one of the following (a, b, or c): a) had systemic CS at least 5 days in past 2 years, b) contraindication to systemic CS, or c) had nasal polyp surgery.</p> <p>CHRONIC IDIOPATHIC URTICARIA (CIU): urticaria more than 6 weeks prior to treatment with Xolair with symptoms present more than 3 days per week despite daily non-sedating H1-antihistamine therapy.</p> <p>IgE-MEDIATED FOOD ALLERGY (all of A, B, C and D): A) baseline IgE greater than or equal to 30 IU/mL, B) positive skin prick test or positive in vitro test for IgE to one or more foods, C) history of allergic reaction that met all of the following (a, b, and c): a) signs and symptoms of a significant systemic allergic reaction, b) reaction occurred within a short period of time following a known ingestion of the food, and c) prescriber deemed this reaction significant enough to require a prescription for an epinephrine auto-injector, and D) patient has been prescribed an epinephrine auto-injector.</p> <p>CONTINUATION THERAPY: ASTHMA: patient responded to therapy and continues to receive an ICS. CRwNP: patient responded after 6 months of therapy and continues intranasal CS. CIU: patient responded to therapy. Exception 1: an exception to the requirement of a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-4/13 therapy (Dupixent) used concomitantly with an ICS.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XOSPATA

Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, FLT3-mutation status
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	AML - approve if the disease is FLT3-mutation positive as detected by an approved test. Lymphoid, Myeloid Neoplasms-approve if the patient has eosinophilia and the disease is FLT3-mutation positive as detected by an approved test.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Lymphoid, Myeloid Neoplasms
Part B Prerequisite	No

XPOVIO

Products Affected

- XPOVIO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Multiple Myeloma-Approve if the patient meets the following (A and B): A) The medication will be taken in combination with dexamethasone AND B) Patient meets one of the following (i, ii, or iii): i. Patient has tried at least four prior regimens for multiple myeloma OR ii. Patient meets both of the following (a and b): a) Patient has tried at least one prior regimen for multiple myeloma AND b) The medication will be taken in combination with bortezomib OR iii. Patient meets both of the following (a and b): a) Patient has tried at least one prior regimen for multiple myeloma AND b) The medication will be taken in combination with Darzalex (daratumumab infusion), Darzalex Faspro (daratumumab and hyaluronidase-fihj injection), or Pomalyst (pomalidomide capsules). Note: Examples include bortezomib/Revlimid (lenalidomide capsules)/dexamethasone, Kyprolis (carfilzomib infusion)/Revlimid/dexamethasone, Darzalex (daratumumab injection)/bortezomib or Kyprolis/dexamethasone, or other regimens containing a proteasome inhibitor, immunomodulatory drug, and/or anti-CD38 monoclonal antibody. Diffuse large B-cell lymphoma Note: this includes patients with histologic transformation of indolent lymphomas to diffuse large B-cell lymphoma)-approve if the patient has been treated with at least two prior systemic therapies.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Treatment of multiple myeloma in combination with daratumumab or pomalidomide
Part B Prerequisite	No

XTANDI

Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Xtandi is being used.
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Prostate cancer-castration-resistant [Metastatic or Non-metastatic] and Prostate cancer-metastatic, castration sensitive-approve if Xtandi will be used concurrently with a gonadotropin-releasing hormone (GnRH) analog [for example: leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets)] or if the patient has had a bilateral orchiectomy. Prostate cancer- Non-Metastatic, Castration-Sensitive - approve if pt has biochemical recurrence and is at high risk for metastasis. [Note: High-risk biochemical recurrence is defined as prostate-specific antigen (PSA) doubling time less than or equal to 9 months.]
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XYREM

Products Affected

- SODIUM OXYBATE (PREFERRED NDCS STARTING WITH 00054)

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with Xywav, Wakix or Sunosi
Required Medical Information	Medication history (as described in Other Criteria field)
Age Restrictions	7 years and older
Prescriber Restrictions	Prescribed by a sleep specialist physician or a Neurologist
Coverage Duration	12 months.
Other Criteria	For Excessive daytime sleepiness (EDS) in patients with narcolepsy, 18 years and older - approve if the patient has tried one CNS stimulant (e.g., methylphenidate, dextroamphetamine), modafinil, or armodafinil and narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT). For EDS in patients with narcolepsy, less than 18 years old-approve if the patient has tried one CNS stimulant (e.g., methylphenidate, dextroamphetamine) or modafinil and narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT). Cataplexy treatment in patients with narcolepsy-approve if narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZEJULA

Products Affected

- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Ovarian, fallopian tube, or primary peritoneal cancer, maintenance therapy - approve if the patient is in complete or partial response after platinum-based chemotherapy regimen and if the patient is new to therapy they must have a trial of Lynparza prior to approval of Zejula. Patients who have had a complete or partial response to first-line platinum based chemotherapy and do not have BRCA altered disease are not required to try Lynparza. Uterine leiomyosarcoma-approve if the patient has BRCA2-altered disease and has tried one systemic regimen. In addition, patients new to therapy must have a trial of Lynparza prior to approval of Zejula. Ovarian, fallopian tube or primary peritoneal cancer in the treatment setting-approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Uterine Leiomyosarcoma, Ovarian, fallopian tube or primary peritoneal cancer-treatment
Part B Prerequisite	No

ZELBORAF

Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	BRAFV600 mutation status required.
Age Restrictions	All diagnoses (except CNS cancer)-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Melanoma, patient new to therapy must have BRAFV600 mutation for approval AND have unresectable, advanced or metastatic melanoma. HCL - must have tried at least one other systemic therapy for hairy cell leukemia OR is unable to tolerate purine analogs and Zelboraf will be used in combination with Gazyva (obinutuzumab intravenous infusion) as initial therapy. Thyroid Cancer-patient has disease that is refractory to radioactive iodine therapy. Erdheim-Chester disease, in patients with BRAF V600 mutation-approve. Central Nervous System Cancer-approve if the patient has BRAF V600 mutation-positive disease AND medication is being used for one of the following situations (i, ii, or iii): i. Adjuvant treatment of one of the following conditions (a, b, c or d): a) Pilocytic astrocytoma OR b) Pleomorphic xanthoastrocytoma OR c) Ganglioglioma/neuroglioma/glioneuronal tumor OR d) pediatric diffuse high-grade glioma OR ii. Recurrent or progressive disease for one of the following conditions (a or b): a) glioma/circumscribed glioma OR b) Glioblastoma OR iii. Melanoma with brain metastases AND the medication will be taken in combination with Cotellic (cobimetinib tablets). Histiocytic Neoplasm-approve if the patient has Langerhans cell histiocytosis and one of the following (i, ii, or iii): i. Multisystem disease OR ii. Pulmonary disease OR iii. Central nervous system lesions AND the patient has BRAF V600-mutation positive disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	Patients with Hairy Cell Leukemia, Non-Small Cell Lung Cancer (NSCLC) with BRAF V600E Mutation, Differentiated thyroid carcinoma (i.e., papillary, follicular, or oncocytic carcinoma) with BRAF-positive disease, Central Nervous System Cancer, Histiocytic Neoplasm
Part B Prerequisite	No

ZEPOSIA

Products Affected

- ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details
Exclusion Criteria	MS-Concurrent use with other disease-modifying agents used for multiple sclerosis. UC- Concurrent Use with a Biologic or with a Targeted Synthetic Disease-modifying Antirheumatic Drug (DMARD) for Ulcerative Colitis
Required Medical Information	Diagnosis
Age Restrictions	UC-18 years and older
Prescriber Restrictions	MS-Prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis. UC-Prescribed by or in consultation with a gastroenterologist
Coverage Duration	1 year
Other Criteria	MS-approve. Ulcerative Colitis, initial-approve if the patient has tried TWO of the following: a preferred adalimumab product, a preferred ustekinumab product, a preferred infliximab product, Rinvoq, Skyrizi, Tremfya. Note-a trial of Simponi SC, Entyvio IV/SC, Omvoh IV/SC, a non-preferred adalimumab product, a non-preferred ustekinumab product or a non-preferred infliximab would also count). Cont tx-approve if the patient has been established on Zeposia. Please Note: preferred adalimumab products include Humira (NDCs starting with -00074), Cyltezo, Yuflyma. Preferred infliximab products include Remicade, Zymfentra. Preferred ustekinumab products include Stelara, Selarsdi, Yesintek.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZOLINZA

Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Cutaneous T-Cell Lymphoma including Mycosis Fungoides/Sezary Syndrome-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZTALMY

Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	2 years and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder-approve if the patient has a molecularly confirmed pathogenic or likely pathogenic mutation in the CDKL5 gene and patient has tried or is concomitantly receiving two other antiepileptic drugs.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZURZUVAE

Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG,
25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	Previous treatment with Zurzuvae during the current episode of postpartum depression
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist or an obstetrician-gynecologist
Coverage Duration	14 days
Other Criteria	Postpartum depression-approve if the patient meets the following (A, B and C): A. Patient meets BOTH of the following (i and ii): i. Patient has been diagnosed with severe depression, AND ii. Symptom onset began during the third trimester of pregnancy or up to 4 weeks post-delivery, AND B. Patient is less than or equal to 12 months postpartum, AND C. Patient is not currently pregnant.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZYDELIG

Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	CLL/SLL-approve if the patient has tried at least one systemic regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	small lymphocytic lymphoma
Part B Prerequisite	No

ZYKADIA

Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Erdheim-Chester Disease-approve if the patient has anaplastic lymphoma kinase (ALK) rearrangement/fusion-positive disease. NSCLC, ALK positive-approve if the patient has advanced or metastatic disease that is ALK positive as detected by an approved test and for patients new to therapy must have a trial of Alecensa or Lorbrena prior to approval of Zykadia. NSCLC, ROS1 Rearrangement-approve if the patient has advanced or metastatic disease. Peripheral T-Cell Lymphoma- approve if patient has ALK-positive anaplastic large cell lymphoma (ALCL).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Inflammatory Myofibroblastic Tumor (IMT) with ALK Translocation. Patients with NSCLC with ROS1 Rearrangement. Erdheim-Chester disease. Peripheral T-Cell Lymphoma.
Part B Prerequisite	No

ZYMFENTRA

Products Affected

- ZYMFENTRA

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a targeted synthetic disease-modifying antirheumatic drug (DMARD).
Required Medical Information	Diagnosis
Age Restrictions	18 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist (initial therapy)
Coverage Duration	Approve through end of plan year
Other Criteria	Crohn's Disease, initial therapy-Approve if the patient is currently receiving infliximab intravenous maintenance therapy or will receive induction dosing with an infliximab intravenous product within 3 months of initiating therapy with Zymfentra. Crohn's Disease, continuation-approve if the patient has had a response to therapy. Ulcerative Colitis, initial therapy-Approve if the patient is currently receiving infliximab intravenous maintenance therapy or will receive induction dosing with an infliximab intravenous product within 3 months of initiating therapy with Zymfentra.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZYTIGA

Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*
- *abirtega*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year
Other Criteria	Prostate Cancer-Metastatic, Castration-Resistant (mCRPC)-Approve if abiraterone is being used in combination with prednisone or dexamethasone and the medication is used concurrently used with a gonadotropin-releasing hormone (GnRH) analog (see Note 1) or the patient has had a bilateral orchiectomy. Prostate cancer-metastatic, castration-sensitive (mCSPC)- approve if the medication is used in combination with prednisone and the medication is concurrently used with a GnRH analog (see Note 1) or the patient has had a bilateral orchiectomy. Prostate Cancer - Regional Risk Group - Approve if the patient meets all of the following criteria (A, B, and C): A) abiraterone is used in combination with prednisone AND B) Patient has regional lymph node metastases and no distant metastases AND C) Patient meets one of the following criteria (i or ii): i. abiraterone with prednisone is used in combination with a GnRH analog (see Note 1) OR ii. Patient has had a bilateral orchiectomy. Prostate cancer-very-high-risk-group-approve if according to the prescriber the patient is in the very-high-risk group, the medication will be used in combination with prednisone, the medication will be used in combination with external beam radiation therapy and the patient meets one of the following criteria (i or ii): i. abiraterone is used in combination with a GnRH analog (see Note 1) OR ii. Patient has had a bilateral orchiectomy. Prostate cancer- radical prostatectomy or post radiation therapy-approve if patient meets (A, B, C and D): A) the medication is used in combination

PA Criteria	Criteria Details
	<p>with prednisone, B) meets (i or ii): i) the patient has prostate specific antigen (PSA) persistence or recurrence following radical prostatectomy or ii) PSA recurrence or positive digital rectal examination (DRE) after radiation therapy, C) patient has pelvic recurrence or positive regional lymph nodes, and D) the medication will be used concurrently with GnRH analog (see Note 1) or the patient has had a bilateral orchiectomy. Salivary Gland Tumors- approve if (A, B and C): A) used in combination with prednisone, B) androgen receptor-positive (AR+) recurrent, unresectable or metastatic tumor, and C) used in combination with a GnRH analog (see Note 1). Note 1: examples of GnRH analogs are leuprolide acetate, Lupron Depot (leuprolide acetate intramuscular injection), Trelstar (triptorelin pamoate intramuscular injection), Zoladex (goserelin acetate subcutaneous implant), Vantas (histrelin acetate subcutaneous implant), Firmagon (degarelix subcutaneous injection), Orgovyx (relugolix tablets).</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Prostate Cancer-Regional Risk Group, Prostate cancer-very-high-risk group, Prostate cancer- radical prostatectomy or post radiation, Salivary Gland Tumors
Part B Prerequisite	No

PART B VERSUS PART D

Products Affected

- *acetylcysteine*
- *acyclovir sodium intravenous solution*
- *albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml*
- *amphotericin b*
- *aprepitant*
- *arformoterol*
- *azathioprine oral tablet 50 mg*
- *BOMYNTRA*
- *budesonide inhalation suspension for nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1 mg/2 ml*
- *CLINIMIX 5%/D15W SULFITE FREE*
- *CLINIMIX 4.25%/D10W SULF FREE*
- *CLINIMIX 4.25%/D5W SULFIT FREE*
- *CLINIMIX 5%-D20W(SULFITE-FREE)*
- *cromolyn inhalation*
- *cyclophosphamide oral capsule*
- *CYCLOPHOSPHAMIDE ORAL TABLET*
- *cyclosporine modified*
- *cyclosporine oral capsule*
- *dronabinol*
- *ENGERIX-B (PF)*
- *ENGERIX-B PEDIATRIC (PF)*
- *ENVARSUS XR*
- *everolimus (immunosuppressive)*
- *formoterol fumarate*
- *gengraf oral capsule*
- *granisetron hcl oral*
- *HEPLISAV-B (PF)*
- *intralipid intravenous emulsion 20 %*
- *ipratropium bromide inhalation*
- *ipratropium-albuterol*
- *JYLAMVO*
- *JYNNEOS (PF)*
- *methotrexate sodium*
- *methotrexate sodium (pf) injection solution*
- *methylprednisolone oral tablet*
- *mycophenolate mofetil*
- *mycophenolate sodium*
- *MYHIBBIN*
- *ondansetron hcl oral solution*
- *ondansetron hcl oral tablet 4 mg, 8 mg*
- *ondansetron oral tablet,disintegrating 4 mg, 8 mg*
- *pentamidine inhalation*
- *PLENAMINE*
- *premasol 10 %*
- *PROGRAF ORAL GRANULES IN PACKET*
- *PULMOZYME*
- *RECOMBIVAX HB (PF)*
- *sirolimus*
- *tacrolimus oral capsule*
- *travasol 10 %*
- *TROPHAMINE 10 %*
- *VARUBI*
- *WYOST*
- *XGEVA*

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index

A

abigale 116
abigale lo 116
abiraterone oral tablet 250 mg, 500 mg . 365,
366
abirtega 365, 366
acetylcysteine 367
ACTEMRA ACTPEN 1, 2
ACTEMRA SUBCUTANEOUS 1, 2
ACTIMMUNE 3
acyclovir sodium intravenous solution ... 367
acyclovir topical ointment 4
ADBRY 7
ADEMPAS 8
AIMOVIG AUTOINJECTOR 9
AKEEGA 10
albuterol sulfate inhalation solution for
nebulization 0.63 mg/3 ml, 1.25 mg/3 ml,
2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml 367
alcohol pads 67
ALECENSA 11, 12
alosetron 13
ALUNBRIG ORAL TABLET 180 MG, 30
MG, 90 MG 15
ALUNBRIG ORAL TABLETS,DOSE
PACK 15
alyq 225
ambrisentan 34
amikacin injection solution 500 mg/2 ml. 16,
17
amphotericin b 367
ampicillin sodium injection recon soln 1
gram, 10 gram 16, 17
ampicillin-sulbactam injection 16, 17
aprepitant 367
ARCALYST 19
arformoterol 367
ARIKAYCE 20
armodafinil 182
ASSURE ID INSULIN SAFETY
SYRINGE 1 ML 29 GAUGE X 1/2 69
AUTGYRO ORAL CAPSULE 160 MG, 40
MG 22

AUSTEDO ORAL TABLET 12 MG, 6 MG,
9 MG 23

AUSTEDO XR ORAL TABLET
EXTENDED RELEASE 24 HR 12 MG,
18 MG, 24 MG, 30 MG, 36 MG, 42 MG,
48 MG, 6 MG 23

AUSTEDO XR TITRATION KT(WK1-4)
ORAL TABLET, EXT REL 24HR DOSE
PACK 12-18-24-30 MG 23
AVMAPKI-FAKZYNJA 24
AVONEX INTRAMUSCULAR PEN
INJECTOR KIT 25

AVONEX INTRAMUSCULAR SYRINGE
KIT 25

AYVAKIT 26
azathioprine oral tablet 50 mg 367
azithromycin intravenous 16, 17
aztreonam 16, 17

B

BALVERSA 27
BELBUCA 165, 166
BELSOMRA 75
BENLYSTA SUBCUTANEOUS 28, 29
benztropine oral 112
BESREMI 30
BETASERON SUBCUTANEOUS KIT .. 31
bexarotene 32, 33
BICILLIN L-A 16, 17
BOMYNTRA 367
BONSITY 293, 294
bosentan oral tablet 34
BOSULIF ORAL CAPSULE 100 MG, 50

MG 35
BOSULIF ORAL TABLET 100 MG, 400

MG, 500 MG 35

BRAFTOVI 36

BRUKINSA ORAL CAPSULE 37

budesonide inhalation suspension for
nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1
mg/2 ml 367

buprenorphine transdermal patch.... 165, 166

C

CABLIVI INJECTION KIT 39

CABOMETYX 40, 41

CALQUENCE (ACALABRUTINIB MAL)	58
.....	42
CAMZYOS	59
.....	43, 44
CAPRELSA ORAL TABLET 100 MG, 300	367
MG	113
carglumic acid	46
CAYSTON	367
cefoxitin	16, 17
ceftazidime	16, 17
cefuroxime sodium injection recon soln 750	367
mg	16, 17
cefuroxime sodium intravenous recon soln	367
1.5 gram	16, 17
CHEMET	48
CIBINQO	49, 50
CIMZIA POWDER FOR RECONST	51, 52
CIMZIA SUBCUTANEOUS SYRINGE	51
KIT 400 MG/2 ML (200 MG/ML X 2)	51, 52
cinacalcet	53
CINRYZE	38
ciprofloxacin in 5 % dextrose intravenous	53
piggyback 200 mg/100 ml	16, 17
clindamycin in 5 % dextrose	16, 17
clindamycin phosphate injection	16, 17
CLINIMIX 5%/D15W SULFITE FREE	367
CLINIMIX 4.25%/D10W SULF FREE	367
CLINIMIX 4.25%/D5W SULFIT FREE	367
CLINIMIX 5%-D20W(SULFITE-FREE)	367
clobazam oral suspension	54
clobazam oral tablet	54
clorazepate dipotassium oral tablet 15 mg,	111
3.75 mg, 7.5 mg	111
colistin (colistimethate na)	16, 17
COMETRIQ ORAL CAPSULE 100	64
MG/DAY(80 MG X1-20 MG X1), 140	64
MG/DAY(80 MG X1-20 MG X3), 60	65
MG/DAY (20 MG X 3/DAY)	66
CONEXXENCE	71
COPIKTRA	111
COSENTYX (2 SYRINGES)	232, 233
COSENTYX PEN (2 PENS)	56
COSENTYX SUBCUTANEOUS	57
SYRINGE 75 MG/0.5 ML	57
COSENTYX UNOREADY PEN	57
COTELLIC	367
CRESEMDA ORAL	367
cromolyn inhalation	367
cyclobenzaprine oral tablet 10 mg, 5 mg	367
cyclophosphamide oral capsule	367
CYCLOPHOSPHAMIDE ORAL TABLET	367
.....	367
cyclosporine modified	367
cyclosporine oral capsule	367
CYLTEZO(CF) PEN	5, 6
CYLTEZO(CF) PEN CROHN'S-UC-HS	5, 6
SUBCUTANEOUS PEN INJECTOR	5, 6
KIT 40 MG/0.8 ML	5, 6
CYLTEZO(CF) PEN PSORIASIS-UV	5, 6
SUBCUTANEOUS PEN INJECTOR	5, 6
KIT 40 MG/0.8 ML	5, 6
CYLTEZO(CF) SUBCUTANEOUS	5, 6
SYRINGE KIT 10 MG/0.2 ML, 20	5, 6
MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8	5, 6
ML	5, 6
CYSTAGON	61
CYSTARAN	60
D	60
dalfampridine	62
DANZITEN	63
dasatinib oral tablet 100 mg, 140 mg, 20 mg,	270
50 mg, 70 mg, 80 mg	270
DAURISMO ORAL TABLET 100 MG, 25	270
MG	64
deferasirox	65
deferiprone	66
DIACOMIT	71
diazepam intensol	111
diazepam oral solution 5 mg/5 ml (1 mg/ml)	111
.....	111
diazepam oral tablet	111
diclofenac sodium topical gel 3 %	265
dimethyl fumarate oral capsule, delayed	265
release(dr/ec) 120 mg, 120 mg (14)-	240
mg (46), 240 mg	72
DOPTELET (10 TAB PACK)	73
DOPTELET (15 TAB PACK)	73
DOPTELET (30 TAB PACK)	73
dotti	116
doxy-100	16, 17
doxycycline hyalate intravenous	16, 17

dronabinol	367
droxidopa	74
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML.....	76, 77, 78
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML.....	76, 77, 78
E	
ELIGARD	106
ELIGARD (3 MONTH).....	106
ELIGARD (4 MONTH).....	106
ELIGARD (6 MONTH).....	106
eltrombopag olamine	234, 235
EMGALITY PEN	79
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML	79
ENBREL MINI.....	80, 81
ENBREL SUBCUTANEOUS SOLUTION	80, 81
ENBREL SUBCUTANEOUS SYRINGE	80, 81
ENBREL SURECLICK.....	80, 81
ENGERIX-B (PF).....	367
ENGERIX-B PEDIATRIC (PF).....	367
ENVARSUS XR.....	367
EPIDIOLEX.....	83
EPRONTIA.....	303
ERIVEDGE.....	86
ERLEADA ORAL TABLET 240 MG, 60 MG	87
erlotinib oral tablet 100 mg, 150 mg, 25 mg	88
ertapenem.....	16, 17
estradiol oral.....	116
estradiol transdermal patch semiweekly ..	116
estradiol transdermal patch weekly.....	116
estradiol-norethindrone acet.....	116
everolimus (antineoplastic) oral tablet 89, 90	
everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg.....	89, 90
everolimus (immunosuppressive)	367
exenatide subcutaneous pen injector 10 mcg/dose(250 mcg/ml) 2.4 ml, 5 mcg/dose (250 mcg/ml) 1.2 ml	104

F	
FASENRA PEN.....	91, 92
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML	91, 92
fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr	304, 305
fingolimod.....	93
FINTEPLA.....	94
FIRMAGON KIT W DILUENT SYRINGE	95
fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml	18
formoterol fumarate	367
FOTIVDA	96
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG	97
FULPHILA	98
fyavolv	116
G	
gabapentin oral tablet extended release 24 hr 300 mg, 600 mg	107
GATTEX 30-VIAL.....	99
GAUZE PAD TOPICAL BANDAGE 2 X 2	68
GAVRETO	100
gefitinib.....	101
genraf oral capsule	367
gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml.....	16, 17
gentamicin injection.....	16, 17
GILOTRIF	102
glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml	103
glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml	103
glutamine (sickle cell).....	82
GOMEKLI ORAL CAPSULE 1 MG, 2 MG	105
GOMEKLI ORAL TABLET FOR SUSPENSION	105
granisetron hcl oral	367
H	
HEPLISAV-B (PF)	367

HERNEXEOS.....	110
HUMIRA (PREFERRED NDCS	
STARTING WITH 00074)	
SUBCUTANEOUS SYRINGE KIT 40	
MG/0.8 ML	5, 6
HUMIRA PEN (PREFERRED NDCS	
STARTING WITH 00074).....	5, 6
HUMIRA(CF) (PREFERRED NDCS	
STARTING WITH 00074)	
SUBCUTANEOUS SYRINGE KIT 10	
MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4	
ML.....	5, 6
HUMIRA(CF) PEN (PREFERRED NDCS	
NDCS STARTING WITH 00074)	
SUBCUTANEOUS PEN INJECTOR	
KIT 40 MG/0.4 ML, 80 MG/0.8 ML..	5, 6
HUMIRA(CF) PEN CROHNS-UC-HS	
(PREFERRED NDCS NDCS	
STARTING WITH 00074).....	5, 6
HUMIRA(CF) PEN PSOR-UV-ADOL HS	
(PREFERRED NDCS NDCS	
STARTING WITH 00074).....	5, 6
hydromorphone oral tablet extended release	
24 hr	165, 166
hydroxyzine hcl oral tablet.....	114
I	
IBRANCE	117, 118
IBTROZI.....	119
icatibant.....	120
ICLUSIG.....	121
IDHIFA.....	122
imatinib oral tablet 100 mg, 400 mg	123, 124
IMBRUVIDA ORAL CAPSULE 140 MG,	
70 MG	125
IMBRUVIDA ORAL SUSPENSION.....	125
IMBRUVIDA ORAL TABLET 140 MG,	
280 MG, 420 MG.....	125
imipenem-cilastatin.....	16, 17
IMKELDI.....	123, 124
INBRIJA INHALATION CAPSULE,	
W/INHALATION DEVICE	126
INGREZZA.....	127
INGREZZA INITIATION PK(TARDIV)	
.....	127
INGREZZA SPRINKLE	127
INLYTA ORAL TABLET 1 MG, 5 MG	130
INPEFA.....	131
INQOVI	132
INREBIC.....	133
INSULIN SYRINGE-NEEDLE U-100	
SYRINGE 0.3 ML 29 GAUGE, 1 ML	29
GAUGE X 1/2.....	70
intralipid intravenous emulsion 20 %	367
ipratropium bromide inhalation	367
ipratropium-albuterol	367
ITOVEBI ORAL TABLET 3 MG, 9 MG	
.....	134, 135
ivermectin oral tablet 3 mg, 6 mg	136
IWILFIN	138
J	
JAKAFI.....	139, 140
JAYPIRCA ORAL TABLET 100 MG, 50	
MG	141, 142
jinteli	116
JUBBONTI	232, 233
JYLLAMVO	367
JYNNEOS (PF).....	367
K	
KALYDECO.....	144
KERENDIA	145, 146
KESIMPTA PEN	147
KISQALI FEMARA CO-PACK ORAL	
TABLET 400 MG/DAY(200 MG X 2)-	
2.5 MG, 600 MG/DAY(200 MG X 3)-2.5	
MG	148, 149
KISQALI ORAL TABLET 200 MG/DAY	
(200 MG X 1), 400 MG/DAY (200 MG X	
2), 600 MG/DAY (200 MG X 3) 148, 149	
KOSELUGO	151, 152
KRAZATI	153, 154
L	
lapatinib.....	155, 156
LAZCLUZE ORAL TABLET 240 MG, 80	
MG	157
LEDIPASVIR-SOFOSBUVIR	158
lenalidomide.....	159, 160
LENVIMA ORAL CAPSULE 10 MG/DAY	
(10 MG X 1), 12 MG/DAY (4 MG X 3),	
14 MG/DAY(10 MG X 1-4 MG X 1), 18	
MG/DAY (10 MG X 1-4 MG X2), 20	
MG/DAY (10 MG X 2), 24 MG/DAY(10	

MG X 2-4 MG X 1), 4 MG, 8 MG/DAY (4 MG X 2)..... 161, 162
 leuprolide subcutaneous kit..... 106
 levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml..... 16, 17
 lidocaine topical adhesive patch,medicated 5 % 163
 lidocan iii 163
 linezolid in dextrose 5%..... 16, 17
 LIVTENCITY 164
 LONSURF 167
 lorazepam intensol 111
 lorazepam oral tablet 0.5 mg, 1 mg, 2 mg 111
 LORBRENA ORAL TABLET 100 MG, 25 MG 168
 LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG..... 169, 170
 LUPRON DEPOT..... 171, 172
 lyllana..... 116
 LYNNPARZA 173, 174
 LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5) 175
M
 MAVYRET ORAL PELLETS IN PACKET 176
 MAVYRET ORAL TABLET..... 176
 megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml) 177
 megestrol oral tablet..... 177
 MEKINIST ORAL RECON SOLN 178, 179
 MEKINIST ORAL TABLET 0.5 MG, 2 MG 178, 179
 MEKTOVI 180
 memantine oral capsule,sprinkle,er 24hr 181
 memantine oral solution..... 181
 memantine oral tablet..... 181
 memantine-donepezil..... 181
 meropenem intravenous recon soln 1 gram, 500 mg 16, 17
 methadone oral solution 10 mg/5 ml, 5 mg/5 ml 165, 166
 methadone oral tablet 10 mg, 5 mg. 165, 166
 methotrexate sodium 367

methotrexate sodium (pf) injection solution 367
 methylprednisolone oral tablet..... 367
 metronidazole in nacl (iso-os)..... 16, 17
 metyrosine 224
 mifepristone oral tablet 300 mg 150
 mimvey 116
 modafinil oral tablet 100 mg, 200 mg.... 182
 MODEYSO..... 183
 morphine oral tablet extended release.... 165, 166
 MOUNJARO 104
 moxifloxacin-sod.chloride(iso)..... 16, 17
 mycophenolate mofetil..... 367
 mycophenolate sodium 367
 MYFEMBREE..... 184
 MYHIBBIN 367
N
 nafcillin injection 16, 17
NAMZARIC ORAL CAPSULE,SPRINKLE,ER 24HR 181
 NAYZILAM 185
 NERLYNX 186
 NEXLETOL..... 187, 188
 NEXLIZET 189, 190
 nilotinib hcl oral capsule 150 mg, 200 mg, 50 mg 288, 289
 nilutamide 191
 NINLARO..... 192
 nitisinone..... 193
 NIVESTYM 194, 195
 norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg 116
 NUBEQA 198
NUCALA SUBCUTANEOUS AUTO-Injector 199, 200
NUCALA SUBCUTANEOUS RECON SOLN 199, 200
NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML 199, 200
 NUEDEXTA 201
 NUPLAZID..... 202
 NURTEC ODT 203
 NYVEPRIA 204
O
 octreotide acetate injection solution..... 205

ODOMZO	206
OFEV	207
OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG.....	208
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	209
OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500 MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6) ..	209
OJJAARA	210
OMNITROPE	108, 109
ondansetron hcl oral solution	367
ondansetron hcl oral tablet 4 mg, 8 mg...	367
ondansetron oral tablet,disintegrating 4 mg, 8 mg	367
ONUREG	211
OPSUMIT	212
OPSYNVI	213
ORENCIA CLICKJECT	214
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML	214
ORGOVYX	215
ORKAMBI ORAL GRANULES IN PACKET	216
ORKAMBI ORAL TABLET	216
ORSERDU ORAL TABLET 345 MG, 86 MG	217
OTEZLA	218
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47).....	218
oxacillin.....	16, 17
oxacillin in dextrose(iso-osm) intravenous piggyback 2 gram/50 ml	16, 17
OXERVATE	219
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG	165, 166
OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)	104

P	
PANRETIN	220
pazopanib	332, 333
PEMAZYRE	221
PEN NEEDLE, DIABETIC NEEDLE 29 GAUGE X 1/2.....	69
penicillamine oral tablet.....	222
PENICILLIN G POT IN DEXTROSE INTRAVENOUS PIGGYBACK 2 MILLION UNIT/50 ML, 3 MILLION UNIT/50 ML	16, 17
penicillin g potassium injection recon soln 20 million unit.....	16, 17
penicillin g sodium.....	16, 17
pentamidine inhalation.....	367
phenobarbital.....	115
pimecrolimus.....	301
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	226
pirfenidone oral capsule	227
pirfenidone oral tablet 267 mg, 801 mg..	227
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	228
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	228
PLENAMINE	367
POMALYST	229
posaconazole oral tablet,delayed release (dr/ec).....	230
premasol 10 %.....	367
PREVYMIS ORAL TABLET	231
PRIVIGEN	137
PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML	84, 85
PROGRAF ORAL GRANULES IN PACKET	367
PROLASTIN-C INTRAVENOUS SOLUTION.....	14
PROLIA	232, 233
PROMACTA	234, 235
promethazine oral.....	114
PULMOZYME	367

pyrimethamine	236
Q	
QINLOCK	237
QULIPTA	238
R	
RADICAVA ORS STARTER KIT SUSP	239
RECOMBIVAX HB (PF)	367
RELEUKO SUBCUTANEOUS	240, 241
REPATHA	242
REPATHA SURECLICK	242
RETACRIT	84, 85
RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG	243
REVLIMID	159, 160
REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG	244
REZDIFRA	245, 246
REZLIDHIA	247
REZUROCK	248
riluzole	249
RINVOQ LQ	252
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG	250, 251
roflumilast	253
ROMVIMZA	254
ROZLYTREK ORAL CAPSULE 100 MG, 200 MG	255
ROZLYTREK ORAL PELLETS IN PACKET	255
RUBRACA	256
rufinamide	257
RYBELSUS	104
RYDAPT	258
S	
sajazir	120
sapropterin	259
SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG	260
SELARSDI SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML	271, 272
SIGNIFOR	261
sildenafil (pulmonary arterial hypertension) oral tablet	225
sirolimus	367
SIRTURO	262
SKYRIZI SUBCUTANEOUS PEN INJECTOR	263
SKYRIZI SUBCUTANEOUS SYRINGE	263
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)	263
SODIUM OXYBATE (PREFERRED NDCS STARTING WITH 00054)	354
sodium phenylbutyrate	223
SOFOSBUVIR-VELPATASVIR	264
SOMAVERT	266
sorafenib	267, 268
SOTYKTU	269
SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG	270
STELARA SUBCUTANEOUS SOLUTION	271, 272
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML	271, 272
STIVARGA	273, 274
STREPTOMYCIN	16, 17
SUCRAID	275
sunitinib malate	276, 277
SYMDEKO	278
SYMLINPEN 120	279
SYMLINPEN 60	279
SYMPAZAN	54
T	
TABRECTA	280
tacrolimus oral capsule	367
tacrolimus topical	301
tadalafil (pulm. hypertension)	225
tadalafil oral tablet 2.5 mg, 5 mg	281
TAFINLAR ORAL CAPSULE	283, 284
TAFINLAR ORAL TABLET FOR SUSPENSION	283, 284
TAGRISSO	285, 286
TALZENNA	287
TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG	288, 289
tazarotene topical cream	290
tazarotene topical gel	290
tazicef injection	16, 17

TAZVERIK.....	291
TEFLARO.....	16, 17
TEPMETKO	292
teriflunomide.....	21
teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)	293, 294
testosterone cypionate.....	128, 129
testosterone enanthate	128, 129
testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)	196, 197
testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram), 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)	196, 197
testosterone transdermal solution in metered pump w/app.....	196, 197
tetrabenazine oral tablet 12.5 mg, 25 mg	295
THALOMID ORAL CAPSULE 100 MG, 50 MG	296, 297
TIBSOVO	298
tigecycline	16, 17
tobramycin in 0.225 % nacl	299
tobramycin inhalation	299
tobramycin sulfate injection solution..	16, 17
tolvaptan.....	300
tolvaptan (polycys kidney dis) oral tablet	143
topiramate oral capsule, sprinkle 15 mg, 25 mg	303
topiramate oral solution	303
topiramate oral tablet	303
torpenz.....	89, 90
travasol 10 %.....	367
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION	306
TREMFYA ONE-PRESS	307, 308
TREMFYA PEN INDUCTION PK- CROHN.....	307, 308
TREMFYA PEN SUBCUTANEOUS PEN INJECTOR 200 MG/2 ML	307, 308
TREMFYA SUBCUTANEOUS SYRINGE	307, 308
tretinoin topical	302
tridacaine ii.....	163
trientine oral capsule 250 mg.....	309, 310
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	311
TRIKAFTA ORAL TABLETS, SEQUENTIAL.....	311
TROPHAMINE 10 %	367
TRULICITY	104
TRUQAP.....	312
TUKYSA ORAL TABLET 150 MG, 50 MG	313
TURALIO ORAL CAPSULE 125 MG..	314
TYENNE AUTOINJECTOR.....	1, 2
TYENNE SUBCUTANEOUS.....	1, 2
U	
UBRELVY	315
UPTRAVI ORAL TABLET	316
UPTRAVI ORAL TABLETS,DOSE PACK	316
V	
VALCHLOR	317
VALTOCO	318
vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg.....	16, 17
vancomycin oral capsule 125 mg, 250 mg	319
VANFLYTA	320
VARUBI	367
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG.....	321, 322
VENCLEXTA STARTING PACK	321, 322
VERZENIO.....	323, 324
vigabatrin	325
vigadroner.....	325
vigpoder	325
VITRAKVI ORAL CAPSULE 100 MG, 25 MG	326
VITRAKVI ORAL SOLUTION	326
VIZIMPRO	327
VONJO	328
VORANIGO ORAL TABLET 10 MG, 40 MG	329
voriconazole.....	18, 330
VOSEVI.....	331
VOWST	334
VUMERTY	335
VYNDAMAX.....	282
VYNDAQEL	282

VYVGART HYTRULO	XTANDI ORAL TABLET 40 MG, 80 MG
SUBCUTANEOUS SYRINGE 336, 337 353
W	
WELIREG.....	338, 339
WYOST	367
X	
XALKORI ORAL CAPSULE.....	340, 341
XALKORI ORAL PELLET 150 MG, 20	
MG, 50 MG.....	340, 341
XDEMVY	342
XELJANZ ORAL SOLUTION	343, 344
XELJANZ ORAL TABLET	343, 344
XELJANZ XR	343, 344
XERMELO	345
XGEVA.....	367
XIFAXAN ORAL TABLET 200 MG, 550	
MG	346, 347
XOLAIR SUBCUTANEOUS AUTO-	
INJECTOR 150 MG/ML, 300 MG/2 ML,	
75 MG/0.5 ML	348, 349
XOLAIR SUBCUTANEOUS RECON	
SOLN	348, 349
XOLAIR SUBCUTANEOUS SYRINGE	
150 MG/ML, 300 MG/2 ML, 75 MG/0.5	
ML.....	348, 349
XOSPATA	350
XPOVIO	351, 352
XTANDI ORAL CAPSULE.....	353
Y	
YESINTEK SUBCUTANEOUS	
SOLUTION.....	271, 272
YESINTEK SUBCUTANEOUS SYRINGE	
45 MG/0.5 ML, 90 MG/ML	271, 272
YUFLYMA(CF) AI CROHN'S-UC-HS. 5, 6	
YUFLYMA(CF) AUTOINJECTOR	
SUBCUTANEOUS AUTO-Injector,	
KIT 40 MG/0.4 ML, 80 MG/0.8 ML..	5, 6
YUFLYMA(CF) SUBCUTANEOUS	
SYRINGE KIT 20 MG/0.2 ML, 40	
MG/0.4 ML	5, 6
Z	
ZEJULA ORAL TABLET.....	355
ZELBORA F.....	356, 357
ZEPOSIA	358
ZEPOSIA STARTER KIT (28-DAY)....	358
ZEPOSIA STARTER PACK (7-DAY)..	358
ZOLINZA	359
ZONISADE.....	303
zonisamide	303
ZTALMY	360
ZURZUVAE ORAL CAPSULE 20 MG, 25	
MG, 30 MG.....	361
ZYDELIG	362
ZYKADIA	363
ZYMFENTRA	364